

FOSFOVIT +

OPTIMIZATION OF FERMENTATION

Mixed mineral nitrogen activator to encourage the multiplication of yeasts for medium nitrogen deficiencies.

↓ OENOLOGICAL USES

Ideal for nitrogen deficiencies when you want to avoid the addition of sulphate.

In fermentation and secondary fermentation the use of **FOSFOVIT+** in the early stages of fermentation makes it possible to give the must ammonium salts and thiamine (0.30%) required for the multiplication and fermentative metabolism.

The cellulose present in **FOSFOVIT+** facilitates the dispersion of the yeast cells within the must to be fermented and distances compounds C8 - C10 (medium chain fatty acids) recognised blastomycetic metabolic inhibitors.

↓ DOSAGE AND INSTRUCTIONS FOR USE

15-20 g/hL

With a dose of 20 g/hL for treatment the current limits set by EEC Regulation 606/2009 for thiamine hydrochloride are reached.

If the objective sought is a sharp increase in the biomass of yeasts, **FOSFOVIT+** should be added at the beginning of the fermentation.

However, for use as a fermentation activator, it is generally recommended to add **FOSFOVIT+** a third of the way through and not at the beginning of alcoholic fermentation, to avoid any phenomenon of excessive yeast multiplication. If you need to use **FOSFOVIT+** at the beginning of the fermentation, it is therefore preferable to associate an organic or complex nutrient.

Dissolve **FOSFOVIT+** in a volume of cold water equal to 3 times its weight. After absorption, mix the must well while racking or pumping over.

↓ PACKAGING AND STORAGE

1 kg and 15 kg bags.

Store in a dry place at room temperature and free of odours. Once you open the bag, the product must be used quickly and cannot be stored.