

# INODOSE 2

## SULFUROUS PRODUCTS

Maximum legally permitted quantities in accordance with Annex 1B of the CE Regulation No. 606/2009.

### ↘ OENOLOGICAL APPLICATIONS

**INODOSE 2** contains potassium metabisulphite releasing 2 g of sulphur dioxide when added to must or wine.

It is particularly useful when low doses of sulphur dioxide are required, for example when wines are being aged in barrel.

### ↘ INSTRUCTIONS FOR USE

The amount of sulphur dioxide required needs to be calculated from the current sulphur dioxide level in the product together with the volume liquid involved.

The tablets should rapidly dissolve. Mix thoroughly.

### ↘ DOSE RATE

One tablet of **INODOSE 2** releases 2g of SO<sub>2</sub>.

When sulfiting, it is necessary to adapt the number of tablets to the quantity of SO<sub>2</sub> desired and the volume of wine to be treated.

Example:

- Volume of wine to be re-sulfited: 2 hL
- Recommended re-adjustment quantity: 2 g/hL
- Number of **INODOSE 2** tablets: 2 tablets.

### ↘ PACKAGING AND STORAGE

**INODOSE 2** is supplied in blister packs containing 48 tablets.

Once the blister has been opened, the tablets should be used immediately.

Store in a dry environment which is well ventilated at a temperature between 5 and 25 °C.

The recommended use by date is marked on the packaging.