

Products for vinification, wine filtration and winery hygiene





Perdomini-IOCAn unrivalled partner



The fact that we operate in all the major wine-producing areas in Italy and around the world testifies to our experience in enhancing wines through improved vinification processes.



We work to further improve our skills daily, engaging in ongoing dialogue with oenologists, research laboratories and wineries – perhaps our best source of knowledge and understanding.



We aim to make winemaking an easier task by helping our clients optimize their processes and time management. This is why we never stop trying to find new formulas that will be more effective and more respectful towards the wines themselves, end consumers and the environment.



Thanks to our hands-on experience and our "Research & Development" work, Perdomini-IOC can offer clients a range of products for vinifying and ageing still and sparkling wines, as well as a complete range of filtration components and products for cleaning and sanitizing wineries.

CONTENTS

- **OPTIMIZATION** OF ALCOHOLIC FERMENTATION
 - Yeast protection
 - 9 Pesticide reduction
 - 10 Detoxify the biomass
 - Optimization of aroma biosynthesis in wines 11
 - 12 Organic nutrition
 - 14 Mixed nutrition
 - Inorganic nutrition

YEASTS FOR **ALCOHOLIC FERMENTATION**

- 19 La Claire range
- IOC range
- 35 Blastosel range

TREATMENTS CONTAINING SELECTED YEAST DERIVATIVES

- 43 Before and during the fermentation
- Aging adjuvants
- 47 Pre-bottling treatments

ENZYMES

- Eno&zymes range
- Zimopec range
- Other enzymes

BACTERIA FOR MALOLACTIC FERMENTATION

- Decision criteria
- Direct-inoculation bacteria
- Bacteria with short re-acclimatization time
- Bacteria with standard re-acclimatization time
- 63 Malolactic fermentation starter
- Products for governing bacterial populations

- 67 Bentonite clays
- Anti-oxidation treatments 69
- 74 Riddling adjuvants
- 76 Alternative solution
- 78 Vegan-friendly chitosan-based fining agents
- Fining agents

STABILIZATION

- Gum arabic: stabilization/protection
- Gum arabic: tartrate stabilization
- Tartrate and calcium stabilization
- Carboxymethylcellulose

TREATMENTS TO CORRECT SPECIFIC **DEFECTS**

- 98 Decolourizing carbon
- 100 Decontaminating carbon
- 101 Anti-reduction treatments
- 101 Acidifiers
- 103 De-acidifiers
- 104 Other acids

105 MICROBIOLOGICAL STABILIZATION

- 105 Sulphites
- 109 Other sulphites

TANNINS

- 111 Limiting antioxidant and antioxydase effects
- 112 Limiting the oxidative degradation of anthocyanins; protecting and stabilizing colour over time
- 115 Special tannin-based products for vinification

126 WOODEN BARREL SUBSTITUTES

127 FeelWood! Range

FILTRATION ADDITIVES

- 130 Perlites
- 131 Fossil shell flours
- 131 Precoat filtration

CLEANING AND SANITIZING

- 135 Acidic detergents
- 136 Alkaline detergents
- 140 Chlorinated alkaline detergents
- 142 Sanitizing detergents
- 143 Eco-friendly sanitizing detergents
- 144 Cleaning filtration equipment and systems
- 145 Other products





Product key



WINES

white I rosé I red I traditional method I charmat methof



APPEARANCE

LIQUID | POWDER / GRANULAR



ORGANIC

CERTIFIED ORGANIC | ALLOWED IN ORGANIC WINEMAKING



PRODUCT THAT CONTAINS AN ALLERGENIC SUBSTANCE (THE ABSENCE OF THIS SYMBOL MEANS THAT THE PRODUCT DOES NOT CONTAIN ALLERGENS)





KOSHER | KOSHER PASSOVER

OPTIMIZATION OF ALCOHOLIC FERMENTATION

Nitrogen is one of the most important nutrients for yeasts. It's a vital element of wine-making since it has a major impact on the outcome of the alcoholic fermentation stage. A lack of nitrogen in the must generally limits the growth of the yeast, leading to problems with fermentation. The type of assimilable nitrogen used (organic or ammoniacal) and the time it's added are two key factors. Numerous scientific studies on the mechanisms connecting yeast nutrition to aroma expression in wines have demonstrated the importance of the nutritional environment surrounding the yeast on the quality of the final product.



YEAST PROTECTION

1 kg 10 kg

ECOBIOL PIED DE CUVE

Targeted fermentation regulator

This product is derived from specially selected yeast strains which, when added to the juice at the pre-multiplication stage, provide a wealth of secondary nutritional factors that allow the yeast cells to stay perfectly functional. When rehydrating active dried yeast, it can be useful to use Ecobiol Pied de Cuve to avoid stuck or sluggish fermentation, as it provides vital micronutrients for the cells so that there is no danger of deficiencies in trophic factors caused by the composition of the must.

Composition: inactive yeast.

Main application: provide sterols and other crucial components to optimize the metabolic activity of the yeast during the rehydration phase.

Dosage: 10-30g/hL of wine.

1 kg 10 kg

ECOBIOL PIED DE CUVE AROM

Targeted fermentation regulator

Specially selected inactive yeast which avoids excess production of volatile acidity and sulphurous odours. Ecobiol Pied de Cuve Arom increases the yeasts' production and expression of aromas in any conditions, even where there is high pressure (sparkling/effervescent wines). Supplying free amino acids to the yeast makes it possible to improve the aroma-production process, leading to more intense, complex wines as early as the end of alcoholic fermentation.

Composition: inactive yeast.

Main application: provide crucial elements for full expression of the sensory potential of the yeast during rehydration.

Dosage: 10-30g/hL of wine.

HYDRA PC

Yeast protector for traditional-method sparkling wines

When added at the rehydration stage, Hydra PC allows yeasts to cope with the difficult conditions they can find when the pied de cuve is being inoculated or at the time of the tirage (SO₂, alcohol, pH, sugar, CO₂ levels, etc.) by reinforcing their cell membranes. Hydra PC is an inactive yeast with a naturally high magnesium content - essential for yeast growth and metabolic activity. When yeasts are growing, a high concentration of magnesium allows their cells to divide more rapidly; when static it allows them to consume sugars more rapidly (through the activation of proton pumps) and to enjoy greater cell membrane stabilization, making them more resistant to tricky conditions.

Composition: inactive yeast.

Main application: optimize the yeast rehydration process.

Dosage: 10-40 g/hL (1: 1 ratio with yeast).

ABSOLUTE MV

Compound to reduce the pesticide levels in must and wine

Product designed to absorb pesticide residues in must and wine. Absolute MV cleanses the must or wine of pesticide residues, optimizes the fermentation kinetics, reduces the latency stage and draws out the aromatic properties of the yeasts.

PESTICIDE REDUCTION

Composition: carbon, yeast cell walls, bentonite clay and yeast protein extract.

Main application: absorb pesticide residues which could compromise the alcoholic fermentation process.

Dosage: 20 - 100 g/hL.

ABSOLUTE SP

Compound to reduce pesticide levels in sparkling wines

Absolute SP yeast hulls absorb any residual pesticides in wines, in particular those which S. cerevisiae shows a strong affinity for. Thanks to this affinity, Absolute SP is ideal for preparing bases to turn into sparkling wines.

Composition: yeast hulls, yeast protein extract.

Main application: absorb pesticide residues which could compromise alcoholic fermentation or secondary fermentation.

Dosage: 20-40g/hL. Legal limit: 40g/hL.

10 kg

1 kg

















DETOXIFICATION

500 g - 5 kg 15 kg

CELLCLEAN

Yeast hulls with a powerful detox action

Using Cellclean encourages smooth alcoholic fermentation and helps limit the risk of sluggish fermentation thanks to its ability to remove toxins and the fact it provides survival factors. In the event of stuck alcoholic or malolactic fermentation, Cellclean cleans up the environment before fermentation is started up again.

Composition: yeast hulls.

Main application: detoxify must.

Dosage:

• For stuck / sluggish fermentation: 20-40g/hL

• As prevention: 10-30g/hL

Legal limit: 40 g/hL.

OENOCELL

Alcoholic fermentation starter and regulator

Inert and with a high level of purity, Oenocell is able to adsorb toxic metals and the toxins produced by any moulds, yeasts and bacteria found in the juice. Thanks to the extent of its fibrous surface, Oenocell provides an important support to yeast cells, allowing them to spread through the juice more easily during fermentation. Oenocell comes in rehydrated form for ease and swiftness of application.

Composition: pre-hydrated α -cellulose fibres.

Main application: adsorb toxic metals and other unwanted substances such as the toxins produced by moulds, yeasts and bacteria in the juice.

Dosage:


- 20-40g/hL during alcoholic fermentation
- 60-80g/hL for must with NTU<50

OPTIMIZATION OF AROMA BIOSYNTHESIS IN WINES

1 kg 5 kg

AROMACTIVIT 182

Targeted nutrients to enhance yeasts' capacity for aroma biosynthesis in wine



- first, a dose of Aromactivit 1, just after adding the yeast, in order to facilitate the creation of a sufficient yeast biomass to ensure a high level of aroma release:
- secondly, a dose of Aromactivit 2 a third of the way through alcoholic fermentation, in order to redirect the metabolic flows towards aroma biosynthesis.

Optimizing the secondary metabolic biosynthesis of yeast using Aromactivit 1&2 leads to higher levels of varietal and fermentative aroma compounds.

Composition:

Aromactivit 1: yeast autolysate, inactive yeast and thiamine.

Aromactivit 2: diammonium phosphate, yeast autolysate and thiamine.

Main application: optimization of secondary metabolic biosynthesis in order to achieve higher levels of varietal and fermentative aroma compounds.

Dosage:

- 1) Just after adding the yeasts: 25-30g/hL of Aromactivit in the must.
- 2) 1/3 of the way through fermentation (after a density loss amounting to 30 points): 15-25g/ hL of Aromactivit 2 to the fermenting must.





OPTIMIZATION OF ALCOHOLIC FERMENTATION


ORGANIC NUTRITION

1 kg 5 kg



ACTIVIT 0

Organic nutrient



Activit O also provides essential minerals and vitamins to yeasts and enhances the expression of fermentative and varietal aromas.

Composition: yeast autolysate.

Main application: nutrition with 100% organic nitrogen to help the alcoholic fermentation stage to draw out all the aromatic potential of the grapes.

Dosage:

10–40g/hL depending on the level of assimilable nitrogen in the must, on the yeast strain used, on the concentration of the sugars to be fermented and on the winemaker's aims for the final product. In general, we recommend adding Activit O at two different times: just after adding the yeast and then 1/3 of the way through alcoholic fermentation.

Assimilable nitrogen deficiency in the must	Recommended nutrition	Completion 1/3 of the way through alcoholic fermentation (density loss: 30-40 points)
Severe	ACTIVIT 0 (20 g/hL)	ACTIVIT 0 (20 g/hL) + DAP
Medium	ACTIVIT O (10 g/hL)	ACTIVIT 0 (10 to 30 g/hL) + DAP
Slight	ACTIVIT O (10 g/hL)	ACTIVIT 0 (10 g/hL) + DAP

1 kg

EXTRA PM

Organic nutrient for secondary fermentation (traditional method)

Extra PM ensures optimal activity on the part of the inoculated yeast and improves its overall wellbeing, especially at the latency stage. As it is an inactive yeast with a high glutathione (GSH) content, it's suitable for governing the fining stage in bottles – it acts effectively to combat reduction during the secondary fermentation stage when using the traditional method. Moreover, it preserves varietal aromas (fruity notes in particular), while adding roundness, elegance and longevity.

Composition: inactive yeast with a high reduced-glutathione (GSH) content.

Main application: activate secondary fermentation when using the traditional method.

Dosage:


• 10-30 g/hL

ORGANIC NUTRITION

1 k

ECOBIOL ICE

Yeast nutrient to enhance freshness sensation of white and rosé wines



Composition: preparation derived from yeast fractions, for the improvement of sensory properties in white and rosé wines.

Main application: Ecobiol Ice enhances the secondary metabolism of the yeast, providing that there is enough nitrogen in the must to ensure fermentative performances. An additional classical nutritive complementation is recommended and should be considered depending on the initial YAN level.

Dosage:

• 30-50 g/hL

1 kg - 5 kg 15 kg

ACTIVIT SAFE ™

Detoxifying nutrient at the 2/3 stage of alcoholic fermentation

Activit Safe TM is a 100% organic nutrient containing a yeast autolysate rich in aminic nitrogen (the form of nitrogen most easily assimilated 2/3 of the way through fermentation) and yeast cell walls that adsorb the inhibitory toxins that can build up during fermentation.

Activit Safe TM both gives a boost to the yeast population – helping them to activate their sugar-metabolizing capacity in the event of slow or sluggish fermentation (or where not enough assimilable nitrogen was available at the early stages) – and limits the stress suffered by the yeast because of an accumulation of toxic substances. Activit Safe TM contains yeast cell walls, which are responsible for adsorbing a number of the short-chain fatty acids produced by the yeast during fermentation that and can have harmful effects on the yeast itself.

Composition: yeast autolysate, yeast hulls.

Main application: in the event of either underestimating the initial level of assimilable nitrogen in the must; over-rapid, sluggish or slow fermentation; overly high or low temperature; overly high alcohol level.

Dosage:

• between 20 and 40g/hL, depending on conditions, when the must arrives at a density of 1020-1010 (the 2/3 point of fermentation): 20g/hL if conditions are difficult; 40g/hL if the fermentation is slowing down or it hasn't been possible to provide adequate nutrition to the yeast. Maximum legal limit 65 g/hL.







MIXED NUTRITION

5 kg 15 kg

5 kg

15 kg

ACTIVIT

Mixed nutrient

Activit helps avoid sluggish alcoholic fermentation or totally stuck fermentation caused by a nitrogen deficiency in the must. Adding Activit provides assimilable nitrogen (ammoniacal nitrogen and amino acids), which is vital for the multiplication of the yeast and keeps them active throughout the fermentation process. Activit also acts as a detoxifier, as it takes out inhibitory fatty acids and puts in sterols and long-chain fatty acids, two elements which increase the resilience of the yeast.

Composition: inactive yeast, diammonium phosphate, thiamine.

Main application: provide ammoniacal nitrogen and amino acid nitrogen in order to ensure an optimal start-up to alcoholic fermentation.

Dosage:

- in must: 20-40g/hL.
- half-way through fermentation: 20-40g/hL (with aeration to facilitate sterol synthesis).

Legal limit: 60 g/hL.

ACTIVIT AD

Mixed nutrient with a high organic-nitrogen content

The main organic base of this targeted nutritional product is a yeast derivative. It provides considerably more amino acids than a deactivated yeast, as well as naturally providing the vitamins and minerals needed by yeasts. Activit AD promotes the production of fermentative aromas thanks to its aminic nitrogen content - an excellent source of fruity and floral esters. Activit AD also limits SO₂ production while increasing the efficiency of sulphuring operations by providing thiamine, a vitamin which helps to limit binding between molecules.

Composition: yeast autolysate, diammonium phosphate, thiamine.

Main application: facilitate the healthy growth of yeast and avoid situations of overpopulation.

Dosage:

• 10-40 g/hL.

Legal limit: 60 g/hL.


MIXED NUTRITION

OPTIMIZATION OF ALCOHOLIC FERMENTATION

10 kg

ECOBIOL

Mixed nutrient



Composition: ammonium sulphate, diammonium phosphate, yeast hulls, α -cellulose fibres.

Main applications: provide essential nutrients for Blastomyces metabolism and remove exogenous and endogenous toxins.

Dosage:

ECOBIOL PERLAGE

Targeted secondary-fermentation regulator

Targeted fermentation regulator for sparkling wine production using either the traditional method or the autoclave. Ecobiol Perlage provides nitrogen - of both mineral and amino acid origin - and vitamins, oligoelements, long-chain fatty acids, sterols and cell-wall polysaccharides that act by adsorbing any inhibitors present (e.g. short-chain fatty acids). This balanced blend encourages a swift fermentation process and has positive effects on aromas and bubble formation, not to mention the considerable advantages in terms of time, process management and money.

Composition: inactive yeast, diammonium phosphate.

Main applications: provide ammoniacal nitrogen and amino acid nitrogen for secondary fermentation.

Dosage: 10-55 g/hL.

- 20g/hL in fermentation and refermentation.
- 50g/hL to treat stuck fermentation.





1 kg



INORGANIC NUTRITION

1 kg 25 kg

PHOSPHATE DIAMMONIQUE

Inorganic nutrient

Composition: diammonium phosphate

Main application: boost yeast multiplication by providing additional nitrogen.

Legal limit: 100 g/hL.










FOSFOVIT

Inorganic nutrient



Composition: diammonium phosphate and thiamine.

Main application: boost yeast multiplication.

Dosage: the usual dose is between 5-40g/hL, depending on the amount of assimilable nitrogen in the must, the yeast strain used and the sugar concentration.

Legal limit: 40 g/hL.

1 kg 15 kg

(K)

1 kg 15 kg

FOSFOVIT+

Inorganic nutrient

Ideal for medium-severity nitrogen deficiencies where the winemaker wishes to avoid adding sulphates. Adding Fosfovit+ to must during fermentation and refermentation (even at the earliest stages) provides the ammonium salts and thiamine needed by the yeast for multiplication and sugar metabolism. The cellulose contained in Fosfovit+ facilitates the even spread of the yeast cells through the biomass being fermented and removes the compounds which inhibit the metabolism of Blastomyces.

Composition: diammonium phosphate, cellulose, thiamine.

Applicazioni: boost yeast multiplication.

Dosage:

• 15-20 g/hL

Legal limit: 20 g/hL.

1 kg 15 kg

(K)

MINVIT

Inorganic nutrient

Ideal for severe nitrogen deficiencies, Minvit can be used in both fermentation and refermentation to provide the ammonium salts and thiamine needed by the yeast for multiplication and fermentation metabolism. The cellulose it contains facilitates the even spread of the yeast cells through the biomass being fermented.

Composition: ammonium sulphate, diammonium phosphate, cellulose, thiamine.

Application: boost yeast multiplication.


• 5-40 g/hL depending on the amount of assimilable nitrogen in the must, the yeast strain used and the sugar concentration.

Legal limit: 40 g/hL.

PHOSPHATES TITRES

Inorganic nutrient for secondary fermentation

This product facilitates the complete break-down of sugar into alcohol by the yeast. Applied at the time of preparing the pied de cuve or when using the traditional method, it boosts the start-up of alcoholic fermentation as well as ensuring a sufficient supply of nitrogen-rich nutrition until fermentation is complete.



Main application: provide nitrogen-rich nutrition for secondary fermentation.

Dosage: 5 g/hL. Legal limit: 6 g/hL.

THIAMINE

Inorganic nutrient

Composition: thiamine hydrochloride (vitamin B1).

Main application: boost the growth of the yeasts in the must and increase their viability during alcoholic fermentation. Acts on carbohydrate metabolism at the time of ketonic acids undergoing decarboxylation to become aldehydes.

Dosage: 0, 05 g/hL. Legal limit: 60 mg/hL.




















YEASTS FOR ALCOHOLIC FERMENTATION

The starter yeasts proposed in this section are the result of a selection process enacted by IOC Group researchers in order to obtain a series of specific results from fermentation.

Each strain has been selected to comply with specific technical objectives, such as the expression of a specific sensory profile, alcohol tolerance, resistance to high temperatures, nitrogen requirements and effect on the colour and softness of the wine.

The great advantage of using selected starter yeasts is that they allow the winemaker to start up fermentation in such a way that their desired outcomes are guaranteed.



LA CLAIRE RANGE



YEASTS FOR RED WINES

500 g

LA CLAIRE EXTREME

Active dry yeast

This yeast has been selected for its ability to start up fermentation in musts with a high sugar content destined to become quality wines with a high alcohol content (Amarone, Primitivo, Grenache...) with only limited volatile acidity production. La Claire eXtreme also shows an outstanding ability to produce elegant, persistent aromas in harmony with the volume created by its prolific production of polysaccharides during fermentation.

Main applications: for musts with a high sugar content destined to become quality wines with a high alcohol content.

Composition: Saccharomyces cerevisiae

Dosage:

- 10-25 g/hL for fermentation
- 30-50 g/hL for stuck fermentation or in the most difficult cases

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
short	from 15 − 35°C	high (17.5% vol.)	low	notes of mature red fruit and jam accompanied by hints of nuts and spices

* upon request at the time of ordering

- 79



Active dry yeast

The distinguishing characteristic of La Claire C58 is the production of polysaccharides and glycerin. La Claire C58 is able to provide wines with highly developed fruity aromas that remain stable over time; in red wines, the total flavonoid count is high (with anthocyanins to the fore), there is good intensity of colour and significant quantities of polysaccharides and glycerin. La Claire C 58 provides wines with greater colour stability right from the early stages of maturation, as well as offering significant improvements in body and softness.



Composition: Saccharomyces cerevisiae

Dosage:

- 10-25 g/hL for fermentation
- 30-50 g/hL in the most difficult cases

RMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
rapid	from 15 - 35°C	high (16% vol.)	low	fruity aromas that remain stable over time

^{*} upon request at the time of ordering

500 g

LA CLAIRE T73

Active dry yeast

La Claire T73 stands out because of its production of fruity notes. In red and rosé wines, it is distinguished by its ability to form a high proportion of total flavonoide – with tannins to the fore – which makes it possible to create wines with a stable intensity of colour. La Claire T73 can be used on a whole range of young red and rosé wines: as well as becoming pleasantly fruity, young and aromatic red wines emerge pleasantly soft and well-balanced. Excellent results can be obtained with grapes such as Sangiovese, Dolcetto, Marzemino and Schiava, but also when making new wines in general.

Main application: for the production of red and rosé wines.

Composition: Saccharomyces cerevisiae

Dosage:

- 10-25 g/hL for fermentation
- 30-50 g/hL in the most difficult cases

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
rapid	from 15 - 35°C	high (17% vol.)	high	enhances fruity notes. Adds softness and balance

* upon request at the time of ordering

YEASTS FOR WHITE AND ROSÉ WINES

LA CLAIRE CGC62

Active dry yeast

La Claire CGC62 is distinguished by its very limited development of volatile acidity, even when vinifying clear musts or those with a high alcohol content. Generous feeding with nitrogen not only guarantees continuous fermentation, but also encourages La Claire CGC62 to produce a significant amount of acetates and even more esters, making it particularly suitable for fresh white wines where a fruity, floral set of aromas is desired.

Main applications: for the production of fresh white wines, rosé wines and nouveau wines, as well as *Charmat*-method sparkling wines.

Composition: Saccharomyces cerevisiae

Dosage:

- 10-25 g/hL for fermentation
- 30-50 g/hL in the most difficult cases

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
rapid	from 15 − 35°C	moderate (14% vol.)	high	maintains and enhances the defining features of the grape variety; enhances fruity, floral and fresh notes.

* upon request at the time of ordering

500 g

LA CLAIRE EM2

Active dry yeast

La Claire EM2 is a varietal strain which is able to free and enhance the terpenic aromatic components peculiar to each grape variety. The aromatic profile of La Claire EM2 is characterized by a positive and intriguing note reminiscent of the reductive notes of Sauvignon Blanc. It also has the right characteristics for producing high-range white wines, including those aged or fermented in barriques, which varietal characteristics are enhanced.

Main application: production of high-range white wines.

Composition: Saccharomyces cerevisiae

Dosage:

- 10–25 g/hL for fermentation
- 30-50 g/hL in the most difficult cases

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
rapid	from 15 - 30°C	high (15% vol.)	average-high	releases varietal terpenic aroma compounds. Greater roundness and softness. Adds reductive notes, enhances fruity and floral notes.













LA CLAIRE VDP

Active dry yeast

The perfect yeast for producing high-quality sparkling wines whether the second fermentation takes place in the bottle or in the autoclave. La Claire VDP makes it possible to obtain fresh white wines endowed with a fruity, floral aromatic structure. Perfect for sparkling white wines, La Claire VDP enhances elegance, fineness, structure and aromatic complexity in the first fermentation, which means that bubble formation in the second fermentation can take place more quickly, cleanly and completely, even when working at low temperatures or with a high alcohol content.

Main applications: secondary fermentation using either the traditional or Charmat method, stuck fermentation.

Composition: Saccharomyces cerevisiae r.f. bayanus

Dosage:

- 10-25 g/hL for fermentation
- 30-50 g/hL in the most difficult cases

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
rapid	from 15 - 32°C	high (15,5% vol.)	average-high	enhances fruity and floral notes

* upon request

at the time of ordering

LA CLAIRE EXTASE

Active dry yeast

Yeast isolated from Sauvignon Blanc grapes in the Bordeaux region. Distinguished by its ability to strongly enhance the strength of aromas - in particular citrussy notes, which provide a very pleasant sense of freshness on the nose and in the mouth. With a proper nitrogen nutrition and by conducting fermentation at low temperatures (16 °C), the development of a fruity aromatic profile (esters-acetates) is favored, while at higher temperatures the development of very intense thiol and citrus aromas is favored. The aromatic intensity provided by La Claire Extase is greater the cleaner the must is (<100 NTU).

Main applications: to enhance the aromatic intensity of white wines.

Composition: Saccharomyces cerevisiae

Dosage:

- 10-25 g/hL for fermentation
- 30-50 g/hL for stuck fermentation or in the most difficult cases

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
rapid	from 13 − 20°C	high (14,5% vol.)	average-high	aromatic intensity, freshness, fruity, thiol, persistence

LA CLAIRE VARIETAL TOUCH

Active dry yeast

Strain isolated in the Marlborough region, New Zealand, La Claire Varietal Touch is characterized by the low production of hydrogen sulphide. La Claire Varietal Touch is the strain par excellence in varietal expression and is capable of recognising the vine variety's typical aromatic notes, thanks to its enzymatic range, which is capable of releasing flavours in the form of precursors. La Claire Varietal Touch is at its very best in the production of important and full-bodied white wines.

Main applications: for the production of varietal white wines and when crafting full-bodied, structured white wines.

Composition: Saccharomyces cerevisiae

Dosage:

- 10-25 g/hL for fermentation
- 30-50 g/hL in the most difficult cases

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
normal	from 14 - 22°C	high (15% vol.)	moderate	expression of aromatic varietal notes

YEASTS FOR SPARKLING WINES

500 g

500 g

LA CLAIRE SP665

Active dry yeast

The cryophilia and alcohol tolerance make La Claire SP665 suitable for the production of high-quality sparkling wines produced both using the traditional method and with re-fermentation in an autoclave. Elegance, fineness, structure and aromatic complexity are enhanced by La Claire SP665 in the first fermentation, which can be followed by guick, clean and complete bubble formation even at low temperatures and with a high alcoholic content.

Main application: production of high-quality sparkling wines both using the traditional method and with re-fermentation in an autoclave.

Composition: Saccharomyces cerevisiae r.f. bayanus

Dosage:

- 10-25 g/hL for fermentation
- 30-50 g/hL for stuck fermentation or in the most difficult cases.

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
rapid	from 10 − 30°C	high (18% vol.)	low	elegance, fineness, structure and aomatic complexity

^{*} upon request at the time of ordering













IOC RANGE



YEASTS FOR BIOPROTECTION



Gaïa



500 g







Metschnikowia fructicola yeast with no fermentation capacity but able to limit the proliferation of unwanted yeasts. Gaïa™ reduces unwanted results and

the risk of alcoholic fermentation starting up too soon. It serves as a way to limit pre-fermentation sulphuring, whether used on must in the fermentation tank or even at earlier stages (on the grapes in the harvesting bins). This yeast also facilitates the inoculation of the selected yeasts (Saccharomyces cerevisiae) introduced at a later stage to govern fermentation.

Another great advantage of this product is that it can be used to protect grape juice during prolonged periods of storage or long-distance transport.

Main application: bioprotection at the pre-fermentation stage.

Dosage:

• 7-20 g/hL. The exact dosage varies according to the time of application and the level of microbial contamination risk (which depends on the timing of operations, temperatures, pH, the level of ripeness of the grapes and the quantity of SO₂ added).

OPTIMAL TEMPERATURE	ALCOHOL	NITROGEN
FOR USE IN MACERATION	TOLERANCE	REQUIREMENTS
$8\text{-}16^{\circ}\text{C}$ (cold pre-fermentation, $8\text{-}12^{\circ}\text{C}$).	very low	low

Gaïa[™] yeast needs the subsequent addition of a selected strain of Saccharomyces cerevisiae in order for alcoholic fermentation to take place.

The temperature of the must is a crucial factor, since the colder the must is, the more Metschnikowia fructicola will dominate over Saccharomyces cerevisiae and the stronger the biocontrol it exerts over an unwanted start to fermentation will be.

TEMPERATURE OF MUST	0°C	8°C	12°C	16°C
Approximate average	Several weeks, or even months, with no fermentation activity	7-10 days or more,	4-5 days, then very	2 days, then very
length of a non-		then very limited	limited fermentation	limited fermentation
fermentation stage		fermentation activity	activity	activity

YEASTS FOR ALCOHOLIC FERMENTATION **IOC RANGE**

Box containing: 10 x 500 g - Gaïa 1 x 16 g - NG-KA

GAÏA™ KIT APPASSIMENTO Low SQLIP

Treatment for use on drying grapes in order to inhibit the growth of Botrytis cinerea

Specially formulated bioprotection treatment for grapes to inhibit the proliferation of Botrytis cinerea while the grapes are drying. The GaïaTM Kit Appassimento is made up of a non-Saccharomyces (Gaïa™) yeast and a specially formulated nutrient (NG-KA) to ensure the right amount of organic nitrogen and vitamins while the grapes are left to dry.


The efficacy of this treatment has been demonstrated through genomic and microbiological analyses, which have highlighted its capacity to prevent or limit the extent of fungal attacks.

Composition: Metschnikowia fructicola yeast (GaïaTM) and a specially formulated nutrient (NG-KA) based on deactivated yeast, diammonium phosphate and thiamine.

Main applications: prevention against attacks from the fungus Botrytis cinerea while grapes are drying prior to being used to craft quality wines.


Dosage:

• To treat 10 tonnes of grapes: 5kg of Kit Gaïa™ yeast and 16g of NG-KA nutrient.



Grapes left to dry are susceptible to Botrytis attacks, especially in the early stages of the drying period, as Botrytis cinerea finds the ideal conditions to proliferate then.

Gaïa™ has been shown to be effective at preventing these infections and maintaining a high level of viability for the whole drying period.



Growth trend of Botrytis cinerea during the grapedrying period.

Data obtained from experiments performed on a large scale following induction of Botrytis growth.

YEASTS NOT PRODUCING SO₂ — IOC BE RANGE

Incapable of producing SO₂, the IOC BE range of yeasts are the result of an innovative selection procedure which uses marker-assisted methods. IOC BE yeasts make it possible to enhance the sensory potential of the product without the side effect of producing SO₂. Most yeasts can release varying amounts of acetaldehyde into wines, and more often than not (although not exclusively), this phenemenon occurs following pre-fermentation sulphuring of the must. As acetaldehyde combines very easily with SO_n in wines, this practice often leads to having to use higher doses of sulphites. On the other hand, thanks to their genetic profile, IOC BE yeasts do not have the capacity to produce large amounts of acetaldehyde, meaning less need for sulphuring.

> 500 g 10 kg



IOC BE-FRUITS LOW SOZ SOLUTIONS

Active dry yeast

A useful tool for revealing fruity esters in white and rosé wines, this product has no SO₂-producing capacity. It can also reduce the formation of acetaldehyde, a molecule which tends to combine with sulphites.

Main application: expression of fruity esters (red fruits, pineapple and citrus fruits) in white and rosé wines.



Composition: Saccharomyces cerevisiae

Dosage:


• 20-30 g/hL

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
from normal to rapid	from 12°C - 24°C for fruity ester expression 12-15°C.	moderate (14 % vol).	low	fruity esters (notes of red fruit, pineapple, citrus fruits)

* upon request at the time of ordering

SO_a production:

Difference in concentration between the added SO, and the total SO, measured in the finished wine (mg/l)



SO₂ formation during the fermentation of two musts according to the yeast inoculated: while the classic yeast produces more SO, than the quantity added initially, in the wines fermented using IOC BE Fruits we note that the sulphite content is lower than it was initially


YEASTS FOR ALCOHOLIC FERMENTATION **IOC RANGE**



IOC BE THIOLS LOW SOZ SOLUTIONS



Active dry yeast



Main application: expression of fruity thiols (citrus and exotic fruits) in white and rosé wines.

Composition: Saccharomyces cerevisiae


Dosage:

• 20-30 g/hL

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
very rapid	from 13 − 25°C. max. thiol expression: 15-18°C	high (15% vol.).	moderate	thiol expression (citrus and exotic fruits), enhancement of 3MH

* upon request at the time of ordering

Total SO₂ concentrations: deviations between wine and must



SO, formation during the fermentation of two musts according to the yeast inoculated: while the classic yeasts produce more SO, than the quantity added initially, in the wines fermented using IOC BE Thiols, the final sulphite level is lower than the initial concentration.









YEASTS FOR ALCOHOLIC FERMENTATION

IOC BE FRESH LOW SOS SOLUTIONS

Active dry yeast

A valuable tool for revealing notes associated with freshness in red wines, this product has no SO₂-producing capacity. It can also reduce the formation of acetaldehyde, a molecule which tends to combine with sulphites.

Main applications: for the production of red wines.

Composition: Saccharomyces cerevisiae


Dosage:

• 20-30 g/hL

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
normal	from 20 − 28 °C	high (15,5% vol.)	high	revelation of aromas associated with fresh fruity notes

Give over-ripe grapes back their freshness, on the nose and the palate

IOC Be Fresh has the ability to release specific aroma compounds which help connote fresh-fruit sensations in red wines. It also makes it possible to increase the concentration of certain straight-chain fatty-acid esters (fresh red fruit) and their branched-chain counterparts (fresh black berries) without losing the unique characteristics of the grape variety.



Moreover, the freshness on the nose is reinforced by the ability of IOC BE Fresh to preserve the malic acid in the grapes, whereas most yeasts tend to consume a part of it during alcoholic fermentation (potential consumption between 10% and 30%).

IOC Be Fresh increases the purity of this freshness because it's incapable of producing sulphites, which are notorious for masking aromas. While most yeasts tend to produce and accumulate sulphites from sulphates - in greater or lesser concentrations depending on the strains and the fermentation conditions - IOC Be Fresh avoids this pitfall. Lastly, with this yeast, the risk of reduction is much lower.

500 g





YEASTS FOR RED WINES

IOC PRIMEROUGE R9001

Active dry yeast

Yeast strain particularly recommended for young red wines and nouveau wines, as it produces rounded, intensely aromatic wines. The right choice when the desired outcome is a pleasant, rounded wine, especially one made using Gamay, Merlot or even Syrah.

Main application: for the production of young red wines and nouveau wines.

Composition: Saccharomyces cerevisiae

Dosage:

• 10-20 g/hL

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
normal	from 14 - 24°C	moderate (14% vol.).	high. To prevent the onset of sulphurous odours, try using complex nutrients.	rounded, intensely aromatic wines with notes of red fruit (strawberry and raspberry), jam and blackcurrant

IOC R-9008

Active dry yeast

Thanks to its production of high levels of polysaccharides, IOC R-9008 makes it possible to limit the sensations caused by acetaldehyde in high-alcohol wines. This is one of the go-to yeasts when aiming for longevity, and it reduces the risks of herbaceous notes and excessive tannicity. In the difficult initial conditions for musts in the hottest vine-growing regions, it helps limit the perception of dryness and bitter notes at the same time as intensifying mineral and salty sensations and increasing the length in the mouth.

Main application: in the production of full-bodied red wines made from grapes with a high sugar concentration harvested when fully ripe.

Composition: Saccharomyces cerevisiae

Dosage:

• 20-30 g/hL

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
normal	from 18 - 30°C	high (16% vol.).	low	expression of ripe and complex fruity aromas, volume in the mouth













IOC FRESH ROSÉ

Active dry yeast

Facilitates the development of floral notes in young wines made from grapes considered as "neutral". The IOC Fresh Rosé strain reveals the fresh varietal characteristics of aromatic grapes (Pinot Noir, Syrah, Cabernet Sauvignon....).

Main application: for the production of rosé wines.

Composition: Saccharomyces cerevisiae

Dosage: • 10-20 g/hL

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
very rapid	from 12 - 25°C	high (16% vol.).	moderate	expression of fresh varietal notes

500 g

IOC REVELATION TERROIR

Active dry yeast

This strain has been chosen from Pinot Noir grapes for its excellent ability to preserve the colour. It increases the colour intensity from 5% to 15% when compared to many other native yeasts. IOC Revelation Terroir is perfect for drawing out fruity notes in Pinot Noir, Gamay, Grenache, Merlot, Carignan and Tempranillo.

Composition: Saccharomyces cerevisiae

Main application: for the production of red wines, to preserve colour.

Dosage:

• 10-30 g/hL

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
normal	from 18 − 30 °C	high (15 % vol.)	high	fruity varietal notes, freshness, fineness and elegance

YEASTS FOR WHITE AND ROSÉ WINES

IOC B 2000

Active dry yeast

IOC B 2000 is particularly recommended when making aromatic white or rosé wines as it assists expression on the nose with grape types and grapes weak in naturally occurring aromatic precursors. It therefore brings out the ester-type aromas whilst also accentuating significant varietal notes in order to provide an intense, fruity and fresh bouquet.

Main application: for the production of aromatic white and rosé wines.

Composition: Saccharomyces cerevisiae

Dosage:

• 10-20 g/hL

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
normal	from 12 - 24°C	moderate (14% vol.)	low	expression of fresh varietal notes

^{*} upon request at the time of ordering









IOC REVELATION THIOLS

Active dry yeast

Yeast specially selected for its ability to express varietal thiols, the aroma compounds responsible for the distinguishing sensory characteristics of several grape varieties, including Sauvignon Blanc, Colombard or Syrah. IOC Revelation Thiols limits the development of vegetal notes in the wine's bouquet.

Main application: for the expression of varietal thiols.

Composition: Saccharomyces cerevisiae

Dosage:

• from 20-30 g/hL

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
normal	Between 15-25°C, ensures smooth fermentation. Generally recommended temperature for thiol expression: 16-18°C.	high (15 % vol.)	moderate	full expression of varietal thiols





IOC TWICE

Active dry yeast


Yeast strain recommended for creating fresh, complex and balanced wines. Particularly suitable for tank fermentation, IOC TwICE has been used most successfully on Chardonnary musts, but is also effective on other varieties (Viognier, Grenache, Semillon, Gros Manseng, etc.).

Main application: for the production of fresh, complex, balanced white wines.

Composition: Saccharomyces cerevisiae

Dosage:

• 20-30 g/hL



* upon request at the time of ordering

500 g

IOC FIZZ +

Active dry yeast

The IOC FIZZ + yeast has been selected to meet the needs of those making highly expressive sparkling wines using the *Charmat* method. It has very good fermentation qualities in secondary fermentation and thus ensures quick and complete secondary fermentation.

Composition: Saccharomyces cerevisiae

Main application: activation of secondary fermentation using the Charmat method.

Dosage:

• 10-20 g/hL

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
normal	from 15 – 35°C	moderate (14% vol.)	moderate	aromatic intensity, fruity notes

YEASTS FOR SPARKLING WINES

500 g 10 kg

Active dry yeast

IOC 18-2007

This strain is highly recommended for use in sparkling wine production using the traditional method, although it can also be used with the *Charmat* method. IOC 18-2007 also makes a very positive contribution to still wines – both white and red – from the following areas: Alsace, Bordeaux, Bourgogne, Provence and the Loire Valley.

Its excellent ability to adapt to the most testing conditions (low pH and low temperature) means that the sugars can be consumed rapidly and completely, thereby avoiding the production of unwanted secondary compounds.

Main applications: for the production of sparkling wines using the traditional method and for the production of still wines – both white and red – in difficult conditions.

Composition: Saccharomyces cerevisiae

Dosage:

white wines: 10-20 g/hLrosé wines: 20-25 g/hL

• secondary fermentation (traditional method): 10-20 g/hL

• to restart stuck fermentation: 20-40 g/hL

MENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
rapid	from 18 − 30 °C	high (15 % vol.)	low	fineness, preservation of the terroir

^{*} upon request at the time of ordering

MULTIFUNCTIONAL YEAST

500 g 10 kg

IOC 11-1002 K

Active dry yeast

Yeast strain which can be used for white, red and rosé wines, recommended for the production of quality wines. The quality of the killer factor in this strain means that it is easily activated and allows it to trigger alcoholic fermentation quickly. IOC 11-1002 K is suitable for use when making high-alcohol wines.

Main application: production of white, red and rosé quality wines.

Composition: Saccharomyces cerevisiae

Dosage:

- da 10-20 g/hL for white and rosé wines
- da 20-25 g/hL for red wines

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
normal	from 12 - 25°C	high (16% vol.)	moderate	expression of the terroir

















BLASTOSEL RANGE

Blastosel


YEASTS FOR RED WINES

YEASTS FOR UNCONVENTIONAL APPLICATIONS

IOC DYNAMIX

Active dry yeast

In partnership with Inter Rhône, our observation and study of the bevahiour of pure yeast cultures taken from different vineyards and blended together has led to the development of IOC Dynamix: a complex blend of five different yeast strains that, thanks to each one's unique features and how they work in synergy, allows winemakers to fully express the potential of the terroir without any worries.



Main application: for the production of red wines.

Composition: Saccharomyces cerevisiae

Dosage:

• 20-30 g/hL

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
normal	from 22°C - 28°C.	high (16 % vol.)	moderate	Wines with clean aromas. (IOC Dynamix avoids spontaneous fermentation start-up and therefore faults in the nose of the wine)

* upon request at the time of ordering

500 g

IOC BIO

Active dry yeast

Certified organic yeast that complies with all the conditions required by EU regulations and by the organic certification body SGS. This means that the yeast does not contain sorbitan monostearate (E491). This yeast also boasts killing power, high alcohol tolerance and adaptability to a wide range of fermentation temperatures.

Main application: for the production of organic red, rosé and white wines.

Composition: Saccharomyces cerevisiae

Dosage:

• 20-30 g/hL

FERMENTATION	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL	NITROGEN	SENSORY
KINETICS		TOLERANCE	REQUIREMENTS	PROFILE
normal	from 14 – 28°C	high (16 % vol.)	low	full and balanced expression of the grape variety and the terroir

* upon request at the time of ordering

500 g





BLASTOSEL DELICE

Active dry yeast

Yeast strain for red wines that releases large quantities of polysaccharides and mannoproteins, making it ideal for complex, high quality wines. Blastosel Delice is distinguished by the complex range of spicy aromas it adds to wines.

Main application: for the production of high quality red wines.

Composition: Saccharomyces cerevisiae

Dosage:

• 15-30 g/hL

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
normal	from 18 - 32°C	high (15,5 % vol.)	average-high	complex range of spicy aromas

BLASTOSEL GRAND CRU

Active dry yeast

This product's distinctive feature is its ability to develop complex aroma compounds with longterm stability and to extract flavonoids, whether in the form of tannins or anthocyanins, leading to long-term colour intensity.

Main application: for the production of full-bodied red wines.

Composition: Saccharomyces cerevisiae r.f. bayanus

Dosage:

- 15-25 g/hL under normal conditions
- 30-40 g/hL for stuck fermentation or in the most difficult cases

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
normal	from 16 − 30°C	high (17,5 % vol.)	low	full expression of complex aroma compounds







500 g

10 kg



10 kg







YEASTS FOR ALCOHOLIC FERMENTATION **BLASTOSEL RANGE**

BLASTOSEL LAMBDA

Active dry yeast

This strain is best suited to making wine from black grapes, when the winemaker wishes to emphasize their varietal notes. Its behaviour towards polyphenols is neutral, so it must be used in tandem with suitable winemaking techniques. Also effective when used for secondary fermentation in the autoclave or when treating stuck fermentation, as it has a strong capacity for fermentation.



500 g

10 kg

500 g

Main applications: for the production of white and rosé wines and for stuck fermentation.

Composition: Saccharomyces cerevisiae r.f. bayanus

Dosage:

• 15-30 g/hL

FERMENTATION	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL	NITROGEN	SENSORY
KINETICS		TOLERANCE	REQUIREMENTS	PROFILE
normal	from 18 − 32°C	high (18% vol.)	average-high	complex range of spicy aromas

YEASTS FOR WHITE AND ROSÉ WINES

BLASTOSEL FR95

Active dry yeast

Strain that displays excellent kinetic activity in fermentation, even at low temperatures or with high alcohol content; the aromatic profile is particularly rich and complex, with strong fruity notes to the fore completed by significant notes of rose.

Main application: for the production of highly drinkable white wines with strong characters and lots of fruity notes.

Composition: Saccharomyces cerevisiae

Dosage:

- 15-25 g/hL under normal conditions
- 30-40 g/hL for stuck fermentation or in the most difficult cases


FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
very rapid	from 12 - 30°C	high (15% vol.).	low	fruity notes, hints of rose

YEASTS FOR ALCOHOLIC FERMENTATION BLASTOSFI RANGE

10 kg

BLASTOSEL WHITE FEEL

Active dry yeast



Main application: for the production of white and rosé wines.

Composition: Saccharomyces cerevisiae

Dosage:

• 10-30 g/hL

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
rapid	from 15 - 24°C	moderate (14,5 % vol.)	low	notes of exotic fruits and fresh fruit

YEASTS FOR SPARKLING WINES

BLASTOSEL DELTA

Active dry yeast

When used on white wines, Blastosel Delta can provide the product with delicate fermentation aromas, even when large volumes are being produced or the must has been clarified. Thanks to its versatility, this strain conserves and draws out the unique varietal characteristics of the

Main applications: for white wine production and to treat stuck fermentation when using the Charmat method.

Composition: Saccharomyces cerevisiae r.f. bayanus

Dosage:

- 15-25 g/hL under normal conditions
- 30-40 g/hL for stuck fermentation or in the most difficult cases.

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
rapid	from 12 - 35°C	high (18% vol.).	low	full and balanced expression of grape variety and terroir





500 g







YEASTS FOR ALCOHOLIC FERMENTATION **BLASTOSEL RANGE**

BLASTOSEL KAPPA

Active dry yeast

This strain is ideal for making still white and rosé wines, bases for sparkling wines and in secondary fermentation. Its production of acetates and esters draws out fresh and fruity notes, and also makes this strain suitable for creating young, drinkable red wines.

Main applications: for the production of white, red and rosé wines, and for secondary and stuck fermentation.

Composition: Saccharomyces cerevisiae

Dosage:

- 15-25 g/hL under normal conditions
- 30-40 g/hL for stuck fermentation or in the most difficult cases

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
rapid	from 13 − 30°C	high (15% vol.)	low	fresh fruity notes

500 g 10 kg

BLASTOSEL P346

Active dry yeast

Yeast recommended for the production of quality white wines. Thanks to its intense beta-lyase and beta-glycosidase activity, it releases intense aromas of fresh fruit (citrus fruits, apple and pear) and white flowers. In fermentation with aroma precursors and organic nitrogen, Blastosel P346 is able to develop a high quantity of aromas evoking white-fleshed and exotic fruits which integrate with the wines's own set of aromas.

Main applications: for the production of aromatic white and rosé wines, secondary fermentation in the autoclave when using the Charmat method.

Composition: Saccharomyces cerevisiae r.f. bayanus

Dosage:

 Bases for sparkling wines: 10-20 g/hL • Secondary fermentation: 10-20 g/hL

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
rapid	from 10 - 18°C	moderate (14,5 % vol.)	average-low	aroma of fresh fruit (citrus fruits, apple and pear), exotic fruits and white flowers

500 g







YEASTS FOR ALCOHOLIC FERMENTATION BLASTOSFI RANGE

MULTIFUNCTIONAL YEAST

500 g 10 kg

BLASTOSEL HORIZON

Active dry yeast

Blastosel Horizon displays good resistance to alcohol, ability to draw out varietal characteristics and clean-cut aromas.

Main application: for the production of white, red and rosé wines.

Composition: Saccharomyces cerevisiae

Dosage:

• 15-30 g/hL

ATION	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL	NITROGEN	SENSORY
CS		TOLERANCE	REQUIREMENTS	PROFILE
اد	from 15 - 30°C	moderate (14,5% vol.)	average-low	enhances aromatic cleanliness and perception of the terroir

BLASTOSEL VS

Active dry yeast

The characteristics of Blastosel VS make it ideal for use in red and rosé wines where lively, intense colour is accompanied by the fruity notes generally found in young, drinkable wines. Its strong alcohol-producing power makes it perfect for use when making high-alcohol wines. Blastosel VS is also suitable for giving a boost to sluggish fermentation and to restart stuck fermentation, especially if the alcohol level is already high.

Main applications: for the production of white, red and rosé wines, secondary and stuck fermentation.

Composition: Saccharomyces cerevisiae r.f. bayanus

Dosage:

- 15-25 g/hL under normal conditions
- 30-40 g/hL in the most difficult cases

FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
normal	from 15 − 30°C	high (15% vol.)	low	fresh fruity and floral notes












Γ	\	
т		

	ТҮРЕ	FERMENTATION	RECOMMENDED	ALCOHOL	NITROGEN	SENSORY
	OF WINE	KINETICS	FERMENTATION TEMPERATURE	TOLERANCE	REQUIREMENTS	PROFILE
LA CLAIRE EXTREME	1	short	from 15 − 35°C	high (17.5% vol.)	low	notes of mature red fruit and jam accompanied by hints of nuts and spices
LA CLAIRE C58		rapid	from 15 - 35°C	high (16% vol.)	low	fruity aromas that remain stable over time
LA CLAIRE T73		rapid	from 15 - 35°C	high (17% vol.)	high	enhances fruity notes. Adds softness and balance
LA CLAIRE CGC62		rapid	from 15 − 35°C	moderate (14% vol.)	high	maintains and enhances the defining features of the grape variety; enhances fruity, floral and fresh notes.
LA CLAIRE EM2		rapid	from 15 - 30°C	high (15% vol.)	average-high	releases varietal terpenic aroma compounds. Greater roundness and softness. Adds reductive notes, enhances fruity and floral notes.
LA CLAIRE VARIETAL TOUCH	Ī	normal	from 14 - 22°C	high (15% vol.)	moderate	expression of aromatic varietal notes
LA CLAIRE SP665	II	rapid	from 10 - 30°C	high (18% vol.)	low	elegance, fineness, structure and aomatic complexity
LA CLAIRE VDP		rapid	from 15 - 32°C	high (15,5% vol.)	average-high	enhances fruity and floral notes
IOC BE-FRUITS		from normal to rapid	from 12°C – 24°C for fruity ester expression 12–15°C.	moderate (14 % vol)	low	fruity esters (notes of red fruit, pineapple, citrus fruits). Avoids aroma masking because of its inability to produce SO ₂
IOC BE THIOLS		very rapid	tra 13 e 25°C. max. thiol expression: 15–18°C	high (15% vol.)	moderate	thiol expression (citrus and exotic fruits), enhancement of 3MH
IOC BE FRESH		normal	from 20 - 28 °C	high (15,5% vol.)	high	revelation of aromas associated with fresh fruity notes
IOC PRIMEROUGE R9001	•	normal	from 14 - 24°C	moderate (14% vol.).	high. To prevent the onset of sulphurous odours, try using complex nutrients.	rounded, intensely aromatic wines with notes of red fruit (strawberry and raspberry), jam and blackcurrant
IOC R-9008	•	very rapid	from 18 - 30°C	high (16% vol.)	low	expression of ripe and complex fruity aromas, volume in the mouth
IOC REVELATION TERROIR	T	normal	from 18 − 30 °C	high (15 % vol.)	high	fruity varietal notes, freshness, fineness and elegance
IOC B 2000		normal	from 12 - 24°C	moderate (14% vol.)	low	expression of fresh varietal notes
IOC FRESH ROSÉ	Ī	very rapid	from 12 - 25°C	high (16% vol.)	moderate	development of floral notes in young wines made from grapes considered "neutral"
IOC REVELATION THIOLS	11	normal	from 15 - 25 °C. for thiol expression: from 16 - 18 °C	high (15 % vol.)	moderate	full expression of varietal thiols

	TYPE OF WINE	FERMENTATION KINETICS	RECOMMENDED FERMENTATION TEMPERATURE	ALCOHOL TOLERANCE	NITROGEN REQUIREMENTS	SENSORY PROFILE
IOC TWICE	1	normal	from 18 e 25 °C. For smooth fermentation, avoid overly high or low temperatures.	high. (15 % vol. when turbidity is greater than 80 NTU)	high	fresh citrus notes (in particular, lemon, peach, apricot and floral notes)
IOC 18-2007		rapid	from 18 - 30 °C	high (15 % vol.)	low	fineness, preservation of the terroir
IOC FIZZ +	1	normal	from 15 - 35°C	moderate (14 % vol.)	moderate	aromatic intensity, fruity notes
IOC 11-1002 K		very rapid	from 12 - 25°C	high (16% vol.)	moderate	expression of the terroir
IOC DYNAMIX	•	normal	from 22°C – 28°C.	high (16 % vol.)	moderate	Wines with clean aromas. (IOC Dynamix avoids spontaneous fermentation start-up and therefore faults in the nose of the wine)
IOC BIO		normal	from 14 - 28°C	high (16 % vol.)	low	full and balanced expression of the grape variety and the terroir
BLASTOSEL DELICE	1	normal	from 18 - 32°C	high (15,5 % vol.)	average-high	complex range of spicy aromas
BLASTOSEL GRAND CRU	1	normal	from 16 - 30°C	high (17,5 % vol.)	low	full expression of complex aroma compounds
BLASTOSEL LAMBDA		normal	from 18 - 32°C	high (18% vol.)	average-high	complex range of spicy aromas
BLASTOSEL FR95	1	very rapid	from 12 - 30°C	high (15% vol.)	low	fruity notes, hints of rose
BLASTOSEL WHITE FEEL	11	rapid	from 15 − 24°C	moderate (14,5 % vol.)	low	notes of exotic fruits and fresh fruit
BLASTOSEL DELTA	11	rapid	from 12 − 35°C	high (18% vol.)	low	full and balanced expression of grape variety and terroir
BLASTOSEL KAPPA		rapid	from 13 - 30°C	high (15% vol.)	low	fresh fruity notes
BLASTOSEL P346		rapid	from 10 - 18°C	moderate (14,5 % vol.)	average-low	aroma of fresh fruit (citrus fruits, apple and pear), exotic fruits and white flowers
BLASTOSEL HORIZON		normal	from 15 - 30°C	moderate (14,5% vol.)	average-low	enhances aromatic cleanliness and perception of the terroir
BLASTOSEL VS		normal	from 15 - 30°C	high (15% vol.)	low	fresh fruity and floral notes

TREATMENTS CONTAINING SELECTED YEAST DERIVATIVES



TREATMENTS CONTAINING SELECTED YEAST DERIVATIVES

BEFORE AND DURING THE FERMENT



GLUTAROM EXTRA LOW SOLUTIONS



Glutathione-rich nutrient

The result of cutting-edge techniques for the selection and production of inactive yeasts with an ultra-high glutathione content. When added at the beginning of fermentation, this product leads to a wine with a much higher GSH content - as long as the yeast is given a sufficient supply of

organic nitrogen. When coupled with low quantities of sulphites, the overall positive impact generated by the high GSH content can be clearly perceived in the sensory profile, even in red wines.

Composition: inactive yeast with a high reduced-glutathione content (GSH>10 mg/g).

Main application: provide reduced glutathione to low SO₂ winemaking processes.

• 15-30 g/hL depending on the must, the grape variety and the sulphite levels.



GLUTAROM

Nutrient with guaranteed glutathione levels

With guaranteed glutathione levels, Glutarom allows the winemaker to preserve fruity notes in young white and rosé wines from an early stage, helping to improve the softness of the wines.

Composition: inactive yeast with guaranteed glutathione levels.

Main application: preserve the freshness of white and rosé wines.

Dosage:

• 15-30 g/hL

Yeast derivatives with guaranteed reduced-glutathione levels

WHAT?	WHEN?	WHY?	GLUTATHIONE CONTENT
GLUTAROM EXTRA	Beginning of alcoholic fermentation	Increases glutathione levels (in white, rosé and red wines) and helps preserve them in the tank and in the bottle, especially in low-sulphite conditions.	**
GLUTAROM	Beginning of alcoholic fermentation	Preserves aromas and enhances softness.	*

1 kg



TREATMENTS CONTAINING SELECTED YEAST DERIVATIVES BEFORE AND DURING THE FERMENTATION

ECOBIOL ROUGE

Inactive yeast for stabilization

When added at the early stages of winemaking, Ecobiol Rouge is effective at stabilizing colour and creating softer, more full-bodied wines. It has undergone a special purification process designed specially with this aim in mind and, when used in tandem with other suitable stabilization techniques such as micro-oxygenation and the addition of exogenous tannins, it leads to more intensely coloured, stable and soft wines with plenty of volume.

Composition: inactive yeast.

Main application: Stabilize colour and mouthfeel (softness and structure).

Dosage:

• 20-30 g/hL of juice.

ECOBIOL SH

Inactive yeast to decrease the reduction smell

A complete organic nutrient whose composition makes it possible to contribute nutritional substances that are indispensable for the correct yeast metabolism and remove light sulphur compounds (hydrogen sulfide and mercaptans) that cause perceptible sensorial deviations during the whole vinification process. Ecobiol-SH makes it possible to reduce the use of copper (metallic tastes), prevent uncontrolled aeration, a possible cause of oxidation, and bring out all of the aromatic qualities of the grape.

Composition: inactive yeast.

Main application: reduce the sulphurous flavours and odours that can develop during alcoholic fermentation.

- 30-40 g/hL on the appearance of a defect for curative action.
- 20 g/hL during alcoholic fermentation for preventive action.



FULLPROTECT LOW SO2 SOLUTIONS



Blend of tannins and yeast-based polysaccharides

Fullprotect is effective thanks to the synergy between certain components of tannins that act to preserve colour, aromas and flavours in white and rosé wines. It particularly targets oxidation in the pre-fermentation stages: the

cell-wall components extracted from a selected deactivated yeast stabilize colour, flavours and aromas, making them less vulnerable to oxidation. The selected tannin extracts, thanks to their level of reactivity to oxygen, limit the oxidation of phenols in the must and reduce the impact of the secondary oxidation mechanisms responsible for the development of unwanted sensory characteristics. Fullprotect is an excellent option for reducing sulphur usage.

Composition: deactivated yeast and tannin extracts.

Main application: protect colour, flavours and aromas in the pre-fermentation stages.

Dosage:

• 30 g/hL

1 kg







1 kg







1 kg



NETAROM EXTRA

Aging adjuvant containing deactivated yeasts

Aging adjuvant containing deactivated yeasts

Netarom Extra can be used in addition to Netarom. Netarom Extra is a copper-rich deactivated yeast selected for its capacity to repress the compounds responsible for the sensation of reduction. It's even effective on wines with significant reduction problems. The polysaccharides released by the formula improve volume and roundness in the mouth. Netarom Extra is a targeted treatment which should be used only in the event of serious faults in the wine.

Composition: deactivated yeast.

Main applications: absorb the sulphurous compounds in wines displaying significant signs of reduction.

Dosage:

• 5-30 g/hL.

NETAROM

in the mouth.


Dosage: • 20-40 g/hL.

Composition: deactivated yeast.



SPHÈRE BLANC

Deactivated yeasts for the aging of white wines



This product is particularly useful when the quality of the fresh lees is low, or when the lees are a source of contamination or sulphurous odours. Sphere Blanc provides an essential contribution to the long-term stabilization of flavours and aromas thanks to the interaction between its polysaccharides and the wine's innate aromas.

Composition: deactivated yeasts.

Main application: stabilize flavours and aromas when aging white wines.

Dosage:

• 10-20 g/hL

AGING ADJUVANTS

Netarom is an aging adjuvant containing deactivated yeasts chosen for their capacity to absorb

the compounds responsible for creating a sense of reduction in the mouth. Through the poly-

saccharides it contains, Netarom improves the volume and complexity of white and red wines

Main application: absorb the sulphurous compounds responsible for the reduction-type smell.















































TREATMENTS CONTAINING SELECTED YEAST DERIVATIVES AGING ADJUVANTS

SPHÈRE EXPRESS

Deactivated yeasts and mannoproteins for aging

Sphere Express is composed of a yeast extract with an extraordinarily high free-mannoprotein content. Also containing peptides and nucleotides, this

veast is particularly effective for the aging process, as it helps increase sensations of volume. fullness and sugar content in a short period of time (from 1 to 8 weeks). This makes Sphere Express an excellent compromise between the deactivated yeasts designed for longer aging periods and the mannoprotein-rich formulas used to produce instantaneous effects shortly before bottling.

Composition: deactivated yeasts and mannoproteins.

Main application: improve volume, length and sugar content when aging wines.

Dosage:

• 5-20 g/hL.

SPHÈRE ROUGE

Deactivated yeasts for the aging of red wines

With Sphere Rouge, the winemaker can quickly increase the immediate perception of volume in the mouth to improve the perception of structure in the

wine. Tannins have a distinct intensity despite manifesting more softly, with the addition of a hint of sweetness at the end. Through the interaction between its polysaccharides and the polyphenols in red wines, Sphere Rouge helps stabilize flavours and aromas in the long term. Moreover, its lack of sulphite reductase activity guarantees its neutrality regarding reductive odours.

Composition: Saccharomyces cerevisiae deactivated yeasts.

Main application: improve volume, length and sugar content when aging red wines.

• 10-20 g/hL.

TREATMENTS CONTAINING SELECTED YEAST DERIVATIVES

PRE-BOTTLING TREATMENTS



1 kg

1 kg


ULTIMA FRESH

Mannoproteins for freshness in the mouth



0,1 kg

1 kg





UltiMA Fresh is a preparation based on selected mannoproteins designed to enhance sensations of freshness in the mouth. Our experiments have demonstrated its ability to rebalance the structure of wines displaying a discrepancy between their phenolic and technological maturity. With an appropriate dosage, UltiMA Fresh reduces bitter and 'green' notes at the same time as increasing freshness in the mouth. When used on less complex wines, it adds to their sweetness and roundness. No matter what wine is being produced, UltiMA Fresh always preserves its originality and unique personality.

Composition: Saccharomyces cerevisiae mannoproteins.

Main application: improve freshness, 'sugariness' and length in the mouth.

Dosage:

• 5-25 g/hL.



ULTIMA SOFT

Mannoproteins for roundness in the mouth



 $0,1 \, \text{kg}$

1 kg



Characteristics: its tendency to contribute to the saltiness of wines and its colloidal action give UltiMA Soft a sensory impact with a lot of potential. Whether used on a white or red wine, the tests we have conducted have demonstrated that even very low doses of UltiMA Soft can shift the balance of acidity and dryness towards roundness, 'sugariness' and aromatic persistence. Nonetheless, the bouquet and the character of the wine are respected and preserved.

Composition: Saccharomyces cerevisiae mannoproteins.

Main application: improve roundess and the persistence of aromas.

Dosage:

• 5-25 g/hL.




ULTIMA READY LIFE

Mannoproteins in solution form for aromatic longevity



1 L



UltiMA Ready Life is a product in solution form which, thanks to the specific mannoprotein extract it contains, interacts positively with the aromatic components in the wine, leading to a longer shelf life. With its ability to increase salinity and provide a colloidal action, ultiMA Ready Life provides some very interesting sensory potential. Even very low doses of UltiMA Ready Life can shift the balance of acidity and dryness towards roundness, 'sugariness' and aromatic persistence. Nonetheless, the bouquet and the character of the wine are respected and preserved.

Composition: Saccharomyces cerevisiae mannoproteins in a solution stabilized with SO_a.

Main application: increase the persistence of aromas.

Dosage:

20-150 mL/hL.











ULTIMA READY EXPRESSION

0,1 kg 1 kg



Mannoproteins in solution form to re-establish freshness and balanced mouthfeel



UltiMA Ready Expression is a product in solution form based on totally soluble mannoproteins which can be added directly to the wine. Thanks to its instant effect, it can be used to treat wine just before bottling. The main effects of using this product can be seen in the aromatic profile, which displays greater freshness and balance. This is particularly useful in the event of a major discrepancy between the phenolic maturity and technological maturity

Composition: Saccharomyces cerevisiae mannoproteins in a solution stabilized with SO_a.

Main applications: improve persistence, increase freshness in the mouth and reduce bitter notes.


Dosage:

• 20-150 mL/hL.










Composition: Saccharomyces cerevisiae mannoproteins in a solution stabilized with SO_a.

Main applications: optimize the mouthfeel of the perlage in wines produced using the Charmat method.

Dosage:

• 5-50 mL/hL.



All the products in the ultiMA range can be microfiltered at 0.45 microns, making them ideal for use just before the final filtration and bottling stages.

1 L





Applications

rosé wines.

• 5-30 mL/hL.

Sensory profile

Contro

Ultima Jump

10 g/hL

Dosage:





Ultima fresh Re-establishes balance.

Rebalances wines with a discrepancy between their phenolic and technological maturity. Increases sweetness and roundness in less complex wines.





ULTIMA JUMP

Composition: Mannoproteins from Saccharomyces cerevisiae.

Mannoproteins for added freshness in white and rosé wines

and reduce astringency and dryness in white and rosé wines. When the right dosage is applied, UltiMA

Jump significantly reduces dryness and astringency at the same time as enhancing sensations of fresh-

ness, roundness and length in the mouth. When used on 'thin' wines, it increases their roundness. No mat-

Main applications: improvement in freshness, roundness and length in the mouth in white and

ter what your treatment needs are, the personality and originality of the wine are always left intact.

100% soluble yeast-derived mannoproteins selected for their ability to increase freshness

Aromatic intensity



Ultima soft Helps achieve balance and softness.

Enhances the saltiness and reduces the dryness of white wines; in red wines, it reduces acidity and astringency and lends greater 'sugariness' and aromatic persistence.







Ultima ready life

Increases the persistence of Gives wines back aromas. their freshness in terms of balanced mouthfeel.

Helps preserve the aromatic profile of white, red and sparkling wines and increases overall persistence.





Freshness

Roundness



ready expression

Performs a targeted

action on red wines to

reduce bitter and astrin-

gent notes: enhances

sapidity in white wines.

Ultima ready fizz Ultima ready expression **Produces** creamier bubbles

Makes a notable difference in obtaining a creamier perlage and improving fineness in sparkling wines produced using the Charmat method.

ulti MA

ready fizz





It contributes to rebalancing the freshness of white wines decreases dryness and astringency conferring softness and aromatic persistence

freshness and

softness.





















100 g 500 g

ENO&ZYMES CLEAR SPEED

Enzymatic preparation for static clarification and flotation

Thanks to its strong pectolytic activity, this enzymatic preparation accelerates the sedimentation of pectin substances in must after pressing, favouring the compaction of the lees even for varieties that are difficult to clarify. Furthermore, thanks to the rapidity of its action, ClearSpeed is suitable for clarification with a flotation unit.

Composition: granules with a high concentration of pectinases, purified by the activity of cinna- myl esterase and anthocyanase. Origin: Aspergillus niger.

Main application: clarification of white and rosé must and wine.

Dosage:

Application: static clarification

• 0,5-2 g/hL - Temperature: 5-12 °C - Time: 3-8 hours.

Application: flotation

• 1 g/hL - Time: at least 2 hours.

100 g 500 g

ENO&ZYMES SWEET PRESS

Enzymatic preparation for maceration on the skins of white grapes

SweetPress favours the release of aromas and aromatic precursors into the must, at the same time reducing the times for maceration on skins/pellicular maceration. Thanks to the secondary enzymatic activities generated by SweetPress, the grape skins become more fragile, which means that less pressure is needed in crushing. This all benefits the quality of the product. The use of SweetPress improves the filterability of wine.

Composition: granules with a high concentration of pectinases, purified by the activity of cinnamyl esterase and anthocyanase. Origin: Aspergillus niger.

Main application: maceration on the skins of white grapes.

Dosage:

• 3 g/q.

FNO&7YMFS RANGE

100 g 500 g

ENO&ZYMES AROM PRESS

Enzymatic preparation for maceration on the skins of white grapes

AromPress favours the release of aromas and aromatic precursors into the must, at the same time reducing the times for maceration on skins. Thanks to the secondary enzymatic activities generated by AromPress, the grape skins become more fragile, which means that less pressure is needed in crushing. This all benefits the quality of the product. The use of AromPress improves the filterability of wine.

Composition: granules with a high concentration of pectinases and beta-glycosidase secondary activities, preventing undesired activities. Origin: Aspergillus niger.

Main applications: maceration on the skins of aromatic white grapes.

• 2-4 g/100 kgs - Temperature: 6-12 °C - Time: 12-14 hours.

500 g

ENO&ZYMES TRUE COLOR

Enzymatic preparation for maceration

TrueColour facilitates the release of anthocyanins into the must, which then remain stable even after aging. As well as showing improved colour, wines treated with this enzyme display a sensory profile which is intenser (red fruit), full-bodied and softer. The use of TrueColour improves the filterability of wine.

Composition: granules with a high concentration of pectinases and specific secondary activities, purified by the activity of cinnamyl esterase and anthocyanase. Origin: Aspergillus niger.

Main applications: maceration of red grapes.

Dosage:

Application: fermentation maceration

• 3-4 g/hL - Temperature: 20-28 °C - Time: 2-8 days.

Application: long fermentation maceration

• 4-5 g/hL - Temperature: 20-28 °C - Time: over 10 days.

Application: pre-fermentation cold maceration

• 2-3 g/hL - Temperature: 8-12 °C Time: 2-4 days.











FN7YMFS FNO&7YMFS RANGE

ENO&ZYMES AROM COLOR

Enzymatic preparation for maceration

AromColor facilitates the optimal and complete breakdown of the colour- and aroma-producing substances contained in grape skins. Thanks to its lack of aggressive secondary activities, AromColor does not encourage the release of astrigent or unpleasant substances. Moreover, its beta-glycosidase activity enhances the expression of aromas right from the earliest stages of winemaking. The use of AromColour improves the filterability of wine.

Composition: granules with a high concentration of pectinases and specific secondary activities (beta-glycosidase), purified by the activity of cinnamyl esterase and anthocyanase. Origin: Aspergillus niger.

Main applications: maceration of red grapes

Dosage:

Application: fermentation maceration

• 3-4 g/hL - Temperature: 20-28 °C - Time: 2-8 days.

Application: long fermentation maceration

• 4-5 g/hL - Temperature: 20-28 °C - Time: più di 10 days.

Application: pre-fermentation cold maceration

• 2-3 g/hL - Temperature: 8-12 °C - Time: 2-4 days.

100 g 500 g

ENO&ZYMES EVOLUTION PLUS

Enzymatic preparation for wine aging

EvolutionPlus accelerates the natural aging processes of wines by encouraging the release of mannoproteins that stabilize unstable colour components. Using EvolutionPlus leads to wines being ready for bottling more quickly and to a reduction in the amount of fining agents needed. The use of EvolutionPlus improves the filterability of wine.

Composition: granules with a high concentration of pectinase, specific secondary activities and β -glucanase $\beta(1-3)$, $\beta(1-6)$, purified by the activity of cinnamyl esterase and anthocyanase. **Origin:** Aspergillus niger and Trichoderma harzianium

Main applications: suitable for use during aging on all kinds of still and sparkling wines; also for treating Botrytis-infected grapes.

Dosage:

Application: when aging white wines

• 2-3 g/hL - Temperature: 8-16 °C - Time: at least 4 weeks.

Application: when aging red wines

• 4-5 g/hL - Temperature: 8-16 °C - Time: at least 4 weeks.

Application: for treatment of Botrytis-infected grapes

• 2-3 g/hL - Temperature: 10-14 °C - Time: 2-4 days.

Application: when aging still/sparkling wines

• 2-4 g/hL - Temperature: 8-12 °C - Time: at least 4 weeks.

100 g 500 g








500 g

ENO&ZYMES PROCESS EXTREME

Enzymatic preparation for static clarification and flotation



Composition: granules with a high concentration of acid pectinase, purified by the activity of cinnamyl esterase and anthocyanase. Origin: Aspergillus niger.

Main applications: clarification of white and rosé wines or maceration of white grapes. Use in difficult conditions with pH < 3 or problematic varieties.

Dosage:

Application: static clarification

• 0,5-2 g/hL - Temperature: 5-12 °C - Time: 3-8 hours.

Application: flotation

• 1 g/hL - Time: 2 hours at least.

500 g









ENO&ZYMES ENZYFLOW

Enzymatic preparation to improve filterability

Thanks to its targeted, synergistic action, EnzyFlow hydrolizes all the molecules (pectins, colloids, glucans, etc.) that can hinder or slow the filtration process, thereby preserving the quality of the must or wine. Using EnzyFlow makes it possible to obtain must and wine that can be filtered at a lower pressure, reduce waste, limit the use of filtration adjuvants, reduce the amount of water needed for rinsing and, above all, increase the filtration yield.

Composition: granules with a high concentration of pectinase, specific secondary activities and β -glucanase $\beta(1-3)$, $\beta(1-6)$, purified by the activity of cinnamyl esterase and anthocyanase. **Origin:** Aspergillus niger **and** Trichoderma harzianium.

Main applications: improve filterability of musts and red, white, still and sparkling wines.

Dosage:

Application: for still wines

• 2-4 g/hL - Temperature: 10-16 °C - Time: at least 48 hours.

Application: for musts

• 3-5 g/hL - Temperature: 10-16 °C - Time: at least 48 hours.

Application: for effervescent/sparkling wines

• 2-4 g/hL - Temperature/Time: secondary fermentation conditions.

ZIMOPEC RANGE LIQUID ENZYMES

1 kg 20 kg

> 1 kg 20 kg

> > 1 kg

20 kg

ZIMOPEC P110L

Enzymatic preparation for static clarification and flotation

Zimopec P110L performs an enzymatic action on soluble and insoluble pectins and helps reduce the viscosity of the product in a short timespan. The advantages of using Zimopec P110L are a greater yield of juice, a reduction in pre-filtration and/or pre-clarification waiting times, greater stability in the final product and a reduction in the quantities of fining agents required.

Composition: enzymatic preparation with pectolytic activity obtained from Aspergillus Niger.

Main application: suitable for the treatment of pressed grapes and must; its ability to completely break down pectins greatly facilitates later fining and filtration operations.

Dosage:

• 0,5-2 mL/hL.

ZIMOPEC PML

Enzymatic preparation for pellicular maceration

Grape pressing is one of the operations in the winemaking process with the strongest influence on both the quality of the final product and total production costs. Zimopec PML has been designed specifically to improve the results of this delicate procedure. The enzymatic action performed on soluble pectins means that the viscosity of the product is quickly reduced but without the skins breaking up at the same time.

Composition: enzymatic preparation with pectolytic activity obtained from Aspergillus Niger.

Main application: targeted product for optimization of grape-pressing operations without an excessive breakdown of solid matter.

Dosage:

• 3-5 mL/q.

ZIMOPEC PXL 09

Enzymatic preparation for maceration

The pectolytic and cellulase activity of Zimopec PXL 09 helps the extraction of colour, both in terms of quantity and quality. Zimopec PXL 09 effectively reduces the pectin content, with the result that the products treated have better filterability and are less susceptible to turbidity phenomena.

Composition: enzymatic preparation with secondary hemicellulase activity obtained from Aspergillus Niger.

Main application: Zimopec PXL 09 is recommended for the treatment of red grapes both in traditional winemaking and modern processes such as Flesh Detente and thermovinification in general.

Dosage:

• 3-5 mL/q.

ZIMOPEC FLOTTOFLASH

Enzymatic preparation for static clarification and flotation

Zimopec Flottoflash is a product developed for flotation. It has enzymatic activity that breaks down pectins and assists the formation and growth of floccules that are easier to separate. The advantages deriving from the use of Zimopec Flottoflash are a reduction in the pre-flotation dwell time, a reduction in the use of clarifiers and better growth of floccules, which is reflected in better clarification.

Composition: enzymatic preparation with pectolytic activity obtained from Aspergillus Niger.

Main application: recommended for treatment during the grape must flotation phase.

Dosage:

• 1-5 mL/hL.

ZIMOPEC 2 FLOTTOFLASH

Enzymatic preparation for static clarification and flotation

Flottoflash 2 is a product developed for flotation in the most difficult cases. It has a high pectolytic activity that breaks down pectins and assists the formation and growth of floccules that are easier to separate. The advantages deriving from the use of Zimopec Flottoflash 2 are a reduction in the pre-flotation dwell time, a reduction in the use of clarifiers and better growth of floccules, which is reflected in better clarification.

Composition: enzymatic preparation with pectolytic activity obtained from Aspergillus Niger.

Main application: recommended for treatment during the grape must flotation phase in the most difficult cases.

Dosage:

• 1-5 mL/hL.

ZIMOPEC CLEAR FLASH

Enzymatic preparation for static clarification and flotation

Clear Flash performs an enzymatic action on soluble and insoluble pectins and helps reduce the viscosity of the product in a short timespan. Moreover, the hydrolysis of pectic substances, which are known to act as protective colloids, means that products treated with Clear Flash display greater filterability.

Composition: enzymatic preparation with pectolytic activity obtained from Aspergillus Niger.

Main application: suitable for the treatment of musts; its ability to completely break down pectins greatly facilitates later fining and filtration operations.

Dosage:

• 1-5 mL/hL.




































ZIMOPEC PRESS FLASH

Enzymatic preparation for pellicular maceration

Performing pellicular maceration on white-wine grapes – especially the more aromatic varieties - has the great disadvantage of making the product difficult to filter because of the large quantity of pectins drawn out. Thanks to its pectolytic action, Zimopec Press Flash makes these kinds of musts notably easier to work with. The enzymatic action performed on soluble pectins lowers the viscosity of the must in a short time, without causing the skins to disintegrate.



Main application: specially designed to enhance grape-pressing operations.

Dosage:

• 3-5 mL/hL.

ZIMOPEC COLOR FLASH

Enzymatic preparation for maceration

The pectolytic action performed by Color Flash makes it possible to lower pectin levels; as a consequence, the treated product displays greater filterability and is less susceptible to clouding and hazing.

Composition: enzymatic preparation with pectolytic activity obtained from Aspergillus Niger.

Main application: suitable for treating grapes destined for use in red wines.

Dosage:

• 3-6 mL/hL.







10 kg







ZIMOPEC RANGE **GRANULAR ENZYMES**

0,5 kg 10 kg







pectin content and the products treated therefore have better filterability and are less susceptible to turbidity phenomena. During the maceration phase, the pectolytic and hemicellulase activity generated by Zimopec PX5 makes it possible to break down the grape cell structure more effectively.

The pectolytic activity generated by Zimopec PX5 translates into a drastic reduction in the

Composition: pectolytic enzymatic preparation with secondary proteolytic and hemicellulase activity, obtained from Aspergillus Niger.

Main applications: suitable for improving wine stability and filterability. When used during maceration, this product helps draw out the aromatic potential and the colour components of the grapes. For use making white or red wines.

Dosage:

• 1-3 g/hL.

ZIMOPEC PX5

Enzymatic preparation to improve filterability

OTHER ENZYMES

IOC EXAROME

Enzymatic preparation to facilitate the release of aroma precursors

As it performs pectinolytic and glucosidic enzymatic actions, Exarome facilitates the release of aroma precursors, which may have been partially glycosilated and therefore "not expressed" during the aging of young wines made from grapes such as Muscat, Gewurztraminer and

Composition: enzymatic preparation with pectolytic and β -glucosidic activity.

Main application: reveal varietal aromas in white wines while aging.




BACTERIA FOR MALOLACTIC FERMENTATION DECISION CRITERIA


BACTERIA FOR MALOLACTIC FERMENTATION

Using specially selected winemaking bacteria allows the winemaker to optimize the malolactic fermentation (MLF) process and avoid indigenous microorganisms in the wine producing aroma-masking molecules. These special bacteria can be inoculated at different stages of the winemaking process, depending on the

conditions that arise and on the winemaker's objectives.

By choosing the right bacterium, re-activation process and inoculation technique (co-inoculation, early inoculation or sequential inoculation), the sensory profile of the wine can be guided towards the desired results.





DIRECT-INOCULATION BACTERIA

25 hL 250 hL

> 25 hL 100 hL

25 hL

250 hL

EXTRAFLORE



Targeted winemaking bacterium

This winemaking bacterium enables malolactic fermentation in white, red and rosé wines. Robust and easy to apply, this bacterium can be incorporated directly into the must or wine without being reactivated first. Extraflore does not produce biogenic amines (this strain does not have the genes responsible for the decarboxylation of amino acids). Its effect on the sensory profile of the wine is to enhance spicy and nutty notes and to lend greater structure.

Composition: Oenococcus Oeni.

Main application: obtain full-bodied, complex wines through co-inoculation or sequential inoculation.

Dosage:

• 1 g/hL.



EXTRAFLORE PURE FRUIT MBR PROCESS



Targeted winemaking bacterium

Targeted bacterium for use in crafting red wines. Acts effectively in difficult conditions for vinification (high alcohol content, low temperatures, excessive acidity or the opposite problem of low malic-acid levels). Its adaptability makes it suitable for use on many different types of wine from different countries. Can be added directly to the must or wine. Thanks to its low diacetyl production and its precise

effect on the sensory profile of the wine, the enzymatic activity of Extraflore Pure Fruit leads to the production of esters which express notes of black and red fruit, enhancing the purity of the fruity sensations in the wine. Wines fermented with Extraflore Pure Fruit also provide a greater feeling of roundness.

Composition: Oenococcus Oeni

Main application: obtain fruity wines through co-inoculation or sequential inoculation.

Dosage:

• 1 g/hL

INOFLORE



Targeted winemaking bacterium

This winemaking bacterium enables malolactic fermentation in white, red and rosé wines. It can be incorporated directly into the must or wine without being reactivated first. Inoflore does not produce biogenic amines (this strain does not have the genes responsible for the decarboxylation of amino acids) and is particularly suited to co-inoculation with yeasts, as this heightens the efficacy of its kinetics. Used with this method, it also produces a very limited amount of diacetyl (buttery notes) and helps reveal and preserve fruity notes, in particular from esters. Undoubtedly the best bacterium to use in co-inoculation and when aiming for a fruity wine.

Composition: Oenococcus Oeni.

Main application: obtain fruity wines through co-inoculation.

Dosage:

• 1 g/hL.

BACTERIA FOR MALOLACTIC FERMENTATION DIRECT-INOCULATION BACTERIA

250 hL

MALIC EXPRESS



Winemaking bacteria specifically developed to carry out very rapid malolactic fermentations. It can be directly incorporated into the wine without rehydrating the product first and is suitable for use under most of the circumstances commonly found in winemaking. Malic-Express does not produce any biogenic amines nor volatile phenols. Inoculate at the end of the alcoholic fermentation stage.

Composition: Oenococcus Oeni.

Main applications: protection of industrial processes and early marketing of red wines.

Dosage:

• 1 g/hL.

BACTERIA WITH SHORT REACCLIMATIZATION TIME

25 hL 100 hL

MAXIFLORE SATINE 13step



Targeted winemaking bacterium

This bacterium is recommended in situations where the grapes being fermented are particularly ripe. It allows the aromatic purity of the wines to be preserved while considerably softening the effects of buttery notes, as it produces only very small quantities of diacetyl (it consumes small quantities of citric acid, and only after the malic acid has been completely used up). In numerous field tests, Maxiflore Satine has also demonstrated its ability to reduce astringent and bitter notes in wines at the same time as enhancing the feeling of volume in the mouth. We recommend using early inoculation (at a density of approx. 1.010), as use of this technique allows the bacterium to exploit the favourable temperature of the wine and to protect red wines from Brettanomyces attacks.

Composition: Oenococcus Oeni bacterium and specially formulated activator containing deactivated yeast.

Main application: reduce astringency and limit buttery notes.

Maxiflore Satine is a kit including:

- a preparation of selected lyophilized lactic bacteria;
- specially formulated activator to add to the must/wine when reacclimatizing the bacteria.

Dosage:

• 1 g/hL.





BACTERIA WITH SHORT RE-ACCLIMATIZATION TIME

25 hL 100 hL

MAXIFLORE ELITE 13step



Targeted winemaking bacterium

Malolactic bacterium which leads to excellent sensory properties in the finished wine. In red wines, it helps enhance the feelings of body and volume in the mouth as well as drawing out spicy notes. In white musts/wines, it can amplify nutty notes, but also protect sensations of freshness (when early inoculation is used). Combining the efficacy of one-step acclimitization with its innate characteristics of resistance to low pH and low temperatures, and to high levels of SO₂ and alcohol, it constitutes a valuable tool for ensuring the success of malolactic fermentation.

Composition: Oenococcus Oeni bacterium and specially formulated activator containing deactivated yeast.

Main application: increase the structure and aromatic complexity of the wine.

Maxiflore Elite is a kit including:

- a preparation of selected lyophilized lactic bacteria;
- specially formulated activator to add to the must/wine when reacclimatizing the bacteria.

Dosage:

• 1 g/hL.

BACTERIA WITH STANDARD REACCLIMATIZATION TIME (REACTIVATION AND ACCLIMATIZATION)

25 hL - 100 hL 1000 hL

INOBACTER

Targeted winemaking bacterium

This winemaking bacterium demonstrates the ability to conduct malolactic fermentation in difficult conditions, in particular where the pH is very low. Requires a reactivation and acclimatization period (pied de cuve).

Composition: Oenococcus Oeni.

Main application: for musts and wines with a very low pH.

Dosage:

• 1 g/hL.


BACTERIA FOR MALOLACTIC FERMENTATION

MALOLACTIC FERMENTATION STARTER

1 kg

NUTRIFLORE FML

MLF optimizer



Nutriflore FML is particularly effective in wines with a low pH (lower than 3.4).

Composition: dectivated yeast.

Main application: provide optimized nutrition in order to speed up malolactic fermentation.


Dosage:

• 20 g/hL.

250 g

NUTRIFLORE PDC

Pied de cuve optimizer




Composition: deactivated yeast.

Main application: nutrient to optimize the malolactic fermentation.

Dosage:

• 25 g/hL (per hL of pied de cuve).







PRODUCTS FOR GOVERNING BACTERIAL POPULATIONS

1 kg

ZIMOPEC OVOLYS

Enzymatic preparation to inhibit lactic bacteria

Ovolys is a valuable tool for governing MLF, when making both red and white wines. Depending on the time of the addition and the dosage applied, Ovolys can stop MLF from taking place, govern its conduction or stabilize the wine once it has reached completion. Ovolys becomes more active as pH raises, and using it as an antibacterial can help reduce the quantity of SO_2 added to the wine.

Composition: Lisozyme hydrochloride.

Main applications: govern malolactic fermentation. Ovolys acts on the latic bacteria (Oenococcus, Leuconostoc, Lactobacillus, Pediococcus) already in the must/wine, but has no effect on the indigenous yeasts or acetic bacteria.

Dosage:

- 25-50 g/hL in white must to prevent start-up of MLF.
- 15-20 g/hL to prevent MLF until alcoholic fermentation is complete.
- Fino a 50 g/hL (max. dose allowed by Reg. CE 1622/00) in white and red wines to halt MLF and stabilize after MLF.





To govern bacterial flora

IOC Sentinel is a totally *allergen-free* and GMO-free product designed to govern the bacterial flora both pre- and post-malolactic fermentation. IOC Sentinel is a new, technological adjuvant made from natural, plant-based ingredients. Its innovative blend of chitin-derived polysaccharides stabilizes red wines after completion of MLF without the onset of the classic diseases, and prevents MLF

from taking place in white and rosé wines where it is not desired. IOC Sentinel also allows Charmat-method base wines to be successfully preserved in low- SO_2 conditions, leading to an optimal start to secondary fermentation. This is an important advantage for winemakers who wish to work with low doses of sulphur. Moreover, IOC Sentinel limits any increases in volatile acidity caused by the acetic bacteria in the wine and has a fining/declouding effect.

Composition: chitosan from Aspergillus niger.

Main application: govern the bacterial flora both pre- and post-malolactic fermentation.

Dosage:

• 25-60 g/hL

PRODUCTS FOR GOVERNING BACTERIAL POPULATIONS

SENTINEL


7

A TRULY INNOVATIVE PRODUCT: NATURAL, BIODEGRADABLE, NON-ALLERGENIC AND GMO-FREE, WITH NO ANIMAL-ORIGIN OR SYNTHETIC SUBSTANCES

Found in many living organisms, chitin is the most plentiful polysaccharide after cellulose. Of all the useful polysaccharides for humans, chitin and its main derivatives (chitosan and chitin-glucan) are becoming more and more popular. In 2003, KitoZyme developed an innovative manufacturing process based on the use of a source of chitin that was not animal-origin but plant-based.

KitoZyme and IOC work in partnership to offer innovative collage products which meet market expectations.


In 2009, only the new technology of manufacturing plant-based chitin derivatives was approved by the Italian Independent Assessment Body; the same products were then authorized for use by the EU in 2010.



7

GOVERNING THE POST-MLF STAGES IN RED WINES


When it comes to red wines, MLF is an important step in the vinification process. Upon completion of MLF, the remaining bacteria can quickly become harmful through increases in pentoses, glycerol and tartaric acid, leading to classic faults in the wine. The most popular treatment used to avoid these problems is sulphuring.



SLUGGISH OR STUCK MLF IN WHITE AND ROSÉ WINES

In white and rosé wines, MLF is quite often not required, as it has a significant effect on the flavour of the product, in particular making it more difficult to attain the desired level of freshness.

In sparkling wines produced using the Charmat method, it's crucial to keep the base wine that will be used for the secondary fermentation intact. The presence of lactic bacteria at this stage is highly risky, as they could compromise the integrity of the batch.



The result of the strategies and instruments developed by IOC to control oxidation and microbiological contamination during the pre-fermentation, fermentation and fining stages, SENTINEL is a precious addition to the winemaker's toolbox as it allows them to reduce SO₂ levels.



4

VINIFICATION AND FINING

Pre- and post-fermentation stages

In order to attain a high level of oxidative resistance, it's vital to act before embarking on the vinification process – after pressing, and before fermentation. Fining musts limits their oxidability and leads to fresher, fruitier profiles in white and rosé wines. This process can be carried out through static fining or flotation. Fining removes the by-products of oxidation which cause browning and can constitute 'aroma traps'. However, during the fining process, not only are the levels of oxidation by-products – orthodiphenols and catechins – reduced, but also the turbidity of the product. It's also an

early colloid-stabilization process which helps keep wine clear over time. Fining brings attractive and repulsive forces into play, in accordance with the electrical charge of the molecules: the potential efficacy of a fining agent depends on its charge density, or in other words on its affinity for the molecules with an opposite charge it needs to remove. The Research & Development team at Perdomini-IOC constantly seeks even better fining solutions using Turbiscan R technology – this allows our experts to monitor the evolution of the clarity of a wine put through a fining process.

Bentonite clays

When exposed to heat, the original grape proteins in white and rosé wines can lead to cloudiness in the bottle. This risk can be successfully avoided by using bentonite clays belonging to the Montmorillonite family. At the end of the alcoholic fermentation process, élevage sur lies reduces protein instability, but the only truly effective treatment to remove the protein aggregates that cause cloudiness is to use bentonite clay. Macro-molecules made up of amino acids and proteins with a positive charge at the pH of the wine will interact with the clay which, when suspended in the liquid, will take on a negative charge. The density of the protein-bentonite particle aggregates is greater than the density of the wine and the flocs precipitate. Targeted testing can be carried out before bottling to establish the risk of protein clouding. The most popular test is the heat test, which involves measuring the difference in turbidity after the wine has been heated to 80°C for 30 minutes. The wine can be considered stable when the delta is less than 2 NTU. From a purely practi-

cal point of view, the greater the clay's capacity to swell in water, the more efficient it will be at removing proteins.

Various types of bentonite clay are available:

Natural bentonite

- Sodium bentonite clays: the cation with the greatest ion-exchange capacity is sodium, a reactive element; the clay swells considerably.
- Calcium bentonite clays: the cation with the greatest ion-exchange capacity is calcium; its swelling and adsorption properties are lesser than the sodium-based clay but it's highly effective at compacting lees.

Activated bentonite

In order to enhance their adsorption capacities, calcium-based clays undergo an activation process using sodium carbonate or sodium hydroxide (depending on the proportion of sodium ions), which are highly reactive and swell considerably. Their immediate effect is equal to or greater than that of sodium bentonite clays, but they show less stability over time.


1 ARIFICATION

BENTONITE CLAYS

1 kg 25 kg

BENTOSTAB GRANULÉS





Composition: activated sodium bentonite.

Main application: clarification of must and wine.

Dosage:

• 20-100 g/hL

BENTOSTAB POUDRE

Activated sodium bentonite

High-efficacy bentonite clay for use in fining and stabilizing must and wine. Displays swift formation of flocs and compact deposits of lees.

Composition: activated sodium bentonite.

Main application: clarification of must and wine.

Dosage:

• 20-100 g/hL

BENT UP

Activated sodium bentonite for flotation

Bent Up is a sodium bentonite clay particularly recommended for flotation with excellent clarification properties. It displays the ability to adsorb proteins and effectively removes unstable protein particles, oxidase enzyme compounds (polyphenol oxidase) and phenol-based particles. In white wines, its capacity to eliminate heat-sensitive proteins avoids the onset of clouding. Strongly recommended for use when the wine has been treated with carbon, as it can remove any residual particles still suspended in the wine.

Composition: activated sodium bentonite.

Main application: clarification of must and wine.

Dosage:

• 10-20 g/hL or more in the most difficult cases




















Activated sodium-calcium bentonite

Inobent is an excellent flocculating agent which facilitates the sedimentation of lees at the end of the alcoholic fermentation process. With low de-proteinizing power, it avoids removing the elements needed for a second fermentation process, therefore is ideal for use in producing sparkling wines. Inobent also contributes to the removal of unstable colour-forming matter, making it suitable for the early stabilization of red wines.



Composition: activated sodium-calcium bentonite.


Dosage:

• 20-100 g/hL

INOBENT NAT

Natural sodium-calcium bentonite

Non-activated bentonite clay for use as a fining agent in must and wine. In granular form for ease of use. Inobent Nat displays excellent declouding properties as it ensures optimal lees sedimentation. Inobent Nat interacts with proteins to provide optimal protein stabilization in white and rosé wines.



Dosage:

• 20-100 g/hL

PENTAGEL

Activated sodium bentonite

Suitable for removing lees from must and for fining and stabilizing wines. When added at the fining stage, Pentagel removes protein compounds, phenolic compounds and part of the pectic substances. When added at the earliest stages of fermentation, it not only performs a clarifying action, but also has a positive impact on the fermentation process.

Composition: activated sodium bentonite.

Main application: clarification of must and wine.

Dosage:

- 20-30 g/hL of must
- 10-20 g/hL of wine.

25 kg

1 kg



1 kg 25 kg





2 kg 10 kg





ANTI-OXIDATION TREATMENTS

1 kg 19.96 kg

PVPP POUDRE

Adsorbent of polyphenolic compounds

PVPP Poudre has the power to eliminate the phenolic substances which cause browning and/ or organoleptic defects in certain wines as well as certain molecules that make musts and wines bitter.

Composition: powder polyvinylpolypyrrolidone.

Main application: clarification of white and red wines.

Dosage:

- 5-10 g/hL of white wine.
- 10-50 g/hL for oxidized red and white wines.

Legal limit: 80g/hL.

1 kg 19.96 kg

PVPP GRANULÉS

Adsorbent of polyphenolic compounds

PVPP Poudre has the power to eliminate the phenolic substances which cause browning and/ or organoleptic defects in certain wines as well as certain molecules that make musts and wines bitter.

Composition: granular polyvinylpolypyrrolidone.

Main application: clarification of white and red wines.

Dosage:

• 20-70 g/hL Legal limit: 80g/hL.

1 kg

METALCLEAN

Fining and stabilizing product containing PVI-PVP

Fining agent with the power to simultaneously eliminate large amounts of metals (it has great affinity in particular with copper) from musts and wines and act on potentially oxidizable molecules, thereby preventing oxidation and general deterioration phenomena from occurring both through direct action and indirect effects, i.e. when metals are removed, they no longer catalyse reactions.

Composition: PVI-PVP, yeast hulls, chitin derivatives from Aspergillus niger.

Main applications: lower the levels of copper and other metals, prevent oxidation responses and pinking effect occurring in white wines at risk.

Dosage:

• 30-80 g/hL

ANTI-OXIDATION TREATMENT

PVI/PVP

Fining and stabilizing product

Polyvinylimidazole and polyvinylpyrrolidone copolymer to reduce the levels of heavy metals in must and wine and to prevent oxidation responses.

Composition: PVI-PVP

Main applications:

- In musts: to reduce the levels of heavy metals (especially copper and iron) and consequently prevent oxidation responses and the risk of cloudiness arising from these.
- In wines: to reduce the levels of heavy metals and to prevent oxidation responses and avoid pinking effect in wines considered at risk.

Dosage:

• 10-50g/hL.

Legal limit: 50 g/hL.

CLEAR GT-F

Composite fining agent for flotation

The association between pea protein, silica gel and PVPP provides for rapid flotation and the effective removal of oxidised and oxidisable phenols, improving the final organoleptic balance. Clear GT-F also acts effectively in reducing the *pinking* phenomenon and stabilises and enhances the fruity flavour. Clear GT-F provides excellent clarification through flotation in a single treatment, leading to a reduction in direct and indirect costs.

Composition: pea protein, silica gel and PVPP.

Main application: rapid and optimal flotation.

Dosage:

• 10-100 g/hL

CLEAR GT-R

Composite fining agent for red must

Clear GT-R allows the winemaker to remove astrigency, vegetal notes and unstable polyphenols without negatively impacting the colour of the wine. It also stabilizes and enhances fruity notes. Must clarification is a fundamental point during the winemaking process and Clear GT-R reduces both the direct and indirect clarification costs in one single treatment.

Composition: targeted yeast hulls, pea protein and silica gel.

Main application: clarification of red must.

Dosage:

• 10-30 g/hL

CLEAR GT-W



1 kg

Composite fining agent for white must

Clear GT-W's clarification action is suitable in preventing oxidative phenomena, *pinking*, protein cloudiness and the formation of bitter flavours in white wines, establishing and enhancing the fruity taste. Must clarification is a fundamental point during the winemaking process and Clear GT-W reduces both the direct and indirect clarification costs in one single treatment.

Composition: PVPP, pea protein and bentonite.

Main application: clarification of white must.

Dosage:

• 10-40 g/hL

P-CELL

Composite fining agent

Composite fining and stabilizing agent and useful fermentation regulator. The fining action of P-Cell is suitable for preventing future oxidation, colour decay protein turbidity and the formation of bitter flavours. Fermentation is the best time to use P-Cell, as in this way it is possible to take advantage of the presence of a-cellulose fibres which function both as support to the yeast cells, as well as adsorbent substances inhibiting yeast metabolisms that may be present in the must.

Composition: PVPP, α -cellulose, activated bentonite.

Main application: clarification of must during fermention.

Dosage:

• 50-120 g/hL

P-FRESH

Composite fining agent

P-Fresh is a clarifier with high fining power that allows simplified filtration, improved colour stability and the elimination of oxidized and oxidisable phenols. Thanks to the presence of activated carbon, P-Fresh is ideal for the clarification of white wines. In addition, P-Fresh combines the qualities of plant proteins with the well-known fining capacity of isinglass, presenting itself as an optimal response to meet the need for a high quality fining agent made from carefully selected raw materials.

Composition: plant proteins, activated vegetable carbon, isinglass.

Main application: clarification of must and wine.

Dosage:

30-60 g/hL

10 kg

15 kg





ANTI-OXIDATION TREATMENTS

P-GREEN

Composite fining agent

When used in must and in white and red wines, P-Green shows a remarkable ability to clarify the liquid, leading to a significant reduction in the oxidized and oxidizable polyphenol content. As it doesn't contain bentonite, the winemaker can deal with protein stabilization separately. Containing plant-based proteins and isinglass, P-Green is the perfect choice for anyone creating allergen-free products and, thanks to our careful selection of base materials, has the necessary characteristics to be used to fine quality wines.

Composition: plant proteins, isinglass.

Main application: clarification of must and wine.

Dosage:

• 20-80 g/hL

P-STAR

Composite fining agent

P-Star effectively combats unstable colloids and also impacts proteins – by combining unstable polyphenol-based particles and making them precipitate, it avoids the onset of browning and the combination of proteins and tannins. The synergistic action of the components in P-Star plays a decisive role in eliminating the risk of future oxidation, protein clouding, pinking and the creation of bitter notes.

Composition: PVPP, activated bentonite, isinglass.

Main application: clarification of must and wine.

Dosage:

• 30-80 g/hL

CASEINATE DE POTASSIUM

Potassium-caseinate-based fining agent

Through an adsorption mechanism, this fining agent is able to react with both polyphenolic substances and metals - iron and copper in particular. The flocs thus formed precipitate quickly and form a compact deposit.

Composition: potassium caseinate.

Main applications: particularly recommended for the prevention and treatment of oxidative faults in wines.

Dosage:

- 50-100 g/hL of must.
- 20-80 g/hL of white wine.
- 10-70 g/hL of red wine.

15 kg







1 kg 20 kg







CASEO CELL

Potassium-caseinate-based fining agent

The optimum balance of proportions between the three active principles makes Caseo Cell an excellent fining agent and a useful fermentation regulator. Caseo Cell is highly soluble and does not leave the product with foreign odours or tastes.

Composition: bentonite, potassium caseinate, α -cellulose.

Main application: clarification of must during fermentation.

Dosage:

• 50–120 g/hL according to the circumstances and operating conditions.

25 kg

CASEO SOL

Potassium-caseinate-based fining agent

Caseo Sol acts on colour keeping it stable over time. Caseo Sol can substitute carbon colour removers when there are slight colour defects to be corrected and it does not strip the wine of its natural aromas. Caseo Sol also acts against metal ions, especially iron and copper, meaning that it can prevent symptoms of chemical and physical instability such as copper casse and ferric phosphate casse.

Composition: potassium caseinate, potassium bicarbonate.

Main application: used to clarify white wines, it helps to attain prolonged stability without affecting the organoleptic properties of the product.

Dosage:

- 20 g/hL as an additive to metal-removal treatment in conjunction with potassium ferrocyanide.
- 40-60 g/hL as a stabilizer for white wines with overly intense colour.
- 80-100 g/hL as a treatment to combat oxidation phenomena.

1 kg 25 kg

CASEO STAR

Potassium-caseinate-based fining agent

Treating white musts with the Caseo Star fining agent allows you to improve the colour quality of the finished product, prevent oxidization reactions affecting the flavans and avoid the most common forms of instability, such as ferric phosphate casse, copper casse, clouding and protein deposits. Using Caseo Star is particularly helpful in maderized wines, where it re-establishes the colour balance by inducing the oxidized phenols (which cause the colour of the wine to darken) to precipitate.

Composition: potassium caseinate, activated bentonite.

Main applications: Caseo Star has been designed to simultaneously attain both the removal of unstable phenolic substances and a drastic reduction in proteins.

Dosage:

- 30-50 g/hL for still wines, including dessert wines.
- 80–100 g/hL for hard-to-fine must and wine.











































RIDDLING ADJUVANTS


1 L 10 L

1 L

1 L

CLARIFIANT S

Fining and stabilizing agent for riddling



Composition: bentonite-clay-based preparation.


Main application: facilitates the riddling of sparkling wines. Can be used both for traditional and automatic riddling.

Dosage:

• 70-80 mL/hL.

CLARIFIANT XL

Fining and stabilizing agent for riddling



Composition: sodium bentonite, silicate solution

Main applications: attain compact sediment in both manual and automatic riddling operations.

Dosage:

60-80 ml/hL.

INOCLAIR 2

Fining agent for riddling

A blend of specially selected bentonite clays with low deproteinizing power which forms a non-sticking skin-like sediment. The low-viscosity alginate ensures cohesion between the bentonite blend and the yeast. When incorporated into wine, Inoclair 2 totally absorbs the yeast and isolates it from the sides of the bottle.

Composition: bentonite, potassium alginate, SO₂.

Main application: fining agent for riddling sparkling wines made using the traditional method.


Dosage:

• 70-90 ml/hL.

RIDDI ING ADJUVANTS

PHOSPHATES MAZURE

Tirage adjuvant for riddling



Composition: kaolin solution, diammonium phosphate, citric acid, SO₂.

Main application: facilitates the riddling of sparkling wines. Can be used both for traditional and automatic riddling.

Dosage:

• 20-30 ml/hl.


The following dosage is recommended:

- Phosphates Mazure: 20 mL/hL
- Clarifiant S: 60 mL/hL

1 L

SOLUTION ST

Riddling adjuvant



Composition: gallic tannin in solution form, copper sulphate, SO₂.

Main applications: improves the aging power of wines and reinforces the antioxidant action of sulphur dioxide.

Dosage:

• 20-40 ml/hL.











ALTERNATIVE SOLUTION

1 kg 15 kg

INOFINE V

Plant-based protein fining agent

Inofine V is a 100% plant-based protein extracted from peas, selected for its reactivity to oxidized and oxidizable phenolic compounds, its efficacy in flocculating suspended matter and making it precipitate, its sensory qualities and its ease of use. To make it even more effective, it can be used in tandem with a fining agent, such as bentonite clay, silica gel or a tannin. When combined with a special winemaking carbon, the treatment is further enhanced and sedimentation takes place even more rapidly.


Composition: pea protein.


Main application: clarification of must and wine.

Dosage:

- 20-40 g/hL to treat white and rosé must during fining or winemaking, depending on the health of the grapes
- 30-50 g/hL to treat grapes in the press
- 10-20 g/hL in white and rosé must undergoing flotation, or to fine white, rosé and red wines

Treatment of a Grenache rosé must during fining - Dosage of 30g/hl for each modality





ALTERNATIVE SOLU

22 kg 1000 kg

INOFINE V MES









Vegetable-protein-based liquid fining agent

Particularly suitable for the preventive treatment of musts easily oxidable, used in wine, Inofine V MES significantly increases resistance to oxidation, improves the organoleptic qualities, reduces the sensation of bitterness and eliminates vegetal or herbaceous notes of wines subject to oxidase cases. In flotation it guarantees a good separation and compaction of the sediment.

Composition: pea protein in a colloidal solution, tartaric acid, SO₂.

Main applications: static clarification and flotation of white and rosé musts.

Dosage:

• in white and rosé must: from 100 to 800 mL/hL depending on harvest health.

• in flotation: 50-100 mL/hL • in white wines: 50-150 mL/hL • in red wines: 100-200 mL/hL

> 22 kg 1000 kg


EASY'UP











Composition: carbon, pea protein, bentonite, tartaric acid, SO₂.

Main application: flotation of white must.

Dosage:

50–150 mL/hL.





VEGAN-FRIENDLY CHITOSAN-BASED FINING AGENTS

QI RANGE

Innovative solutions from IOC: natural, biodegradable, hypoallergenic and free from products of animal origin. The IOC "Qi" range is the result of combining an exclusive, 100% fungal-origin (Aspergillus niger) chitosan with other carefully selected ingredients.

1 kg 15 kg

1

QI NO [OX]

Fining and stabilizing agent

An alternative to casein, Qi No[OX] is a non-allergenic formula which contains no animal-origin or synthetic ingredients. It has been developed for use as a powerful antioxidant for must and wine. Qi No[OX] is a technological adjuvant composed of chitin derivatives and bentonite clay, which

facilitate swift sedimentation. Can be used on white must displaying a tendency to oxidize, or on wines – even if they've already become oxidized – to eliminate brown shades and notes of caramel and Madeira. Also eliminates vegetal and bitter notes.


Composition: chitosan from Aspergillus niger, bentonite.

Main application: clarification of quality white and red wines.

Dosage:

• Must: 30-80 g/hL

• White or rosé wine: 20-60 g/hL



Qi No(OX) corrects colour faults arising from wine oxidatio


The chelating effect of the plant-origin polysaccharides leads to the elimination of copper and iron, the main catalysts of any form of oxidation.

The test presented below shows the monitoring of the yellow tints in a white wine (OD 420nm), providing incontestable proof of the power of Qi No(OX) to restore colour to oxidized wines.

Chelating effect of Qi No(OX) on iron and copper in a white wine.


VEGAN-FRIENDLY CHITOSAN-BASED FINING AGENTS

DO 420nm of a white wine contaminated with several doses of Cu²⁺ e di Fe³⁺ and treated with several doses of Qi No(OX)





 $\mathsf{No}[\mathsf{0x}]$ correcting the sensory faults linked to wine oxidation





Clarificat

VEGAN-FRIENDLY CHITOSAN-BASED FINING AGENTS



1 kg 15 kg

QI FINE

Fining and stabilizing agent

Qi Fine has been formulated using chitosan and pea protein. The special fungal-origin chitosan contained in Qi Fine displays a high density charge, meaning it enables rapid flocculation and sedimentation. The synergy between the chitosan and a pea protein selected because of its level of reactivity to phenolic compounds leads to a distinct improvement in sensory properties, as bitter

sensations and excessive astringency are eliminated.

Composition: chitosan from Aspergillus niger, pea protein, tartaric acid.


Main application: clarification and stabilization.


Dosage:

• in must: 10-50 g/hL • in wine: 10-30 g/hL



Used for improving red must and wine, either during the racking or alcoholic fermentation stages, Qi FINE is highly effective at absorbing the polyphenolic compounds involved in the oxidation process.






In this experiment, Qi FINE demonstrated its ability to flocculate much more rapidly than other fining agents, with an average transmission value of 40% (clearly visible change), achieved in just 3 hours. Other fining agents take 6-8 hours to reach the same level.

VEGAN-FRIENDLY CHITOSAN-BASED FINING AGENTS

The depth of the sediment correlates to the average transmission value. For **Qi FINE**, a sediment with a depth of 0.6mm is a highly satisfactory sign, as it means the lees have settled well and less wine will be lost. Product P, on the other hand, fines the wine but without the lees settling well (sediment depth of only 0.3mm).





QI FINE MES

Fining and stabilizing agent







Qi Fine Mes is the result of combining chitosan and pea protein in an aqueous solution, making it ready for use. The special fungal-origin chitosan contained in Qi Fine Mes displays a high density charge, meaning it enables rapid flocculation and sedimentation. The synergy between the chitosan and a pea protein selected because of its level of reactivity to phenolic compounds leads to a distinct improvement in sensory properties, as bitter sensations and excessive astringency are eliminated.

Composition: chitosan from Aspergillus niger, pea protein, tartaric acid.

Main applications: fining and flotation.

Dosage:

• 50-300 mL/hL





VEGAN-FRIENDLY CHITOSAN-BASED FINING AGENTS



Flotation consists of separating the particles suspended in the must by injecting a gas for them to attach to. This process requires the use of an adjuvant to enable the formation of flocs by encouraging particles and pectins to agglomerate to each other. It's also useful to add enzymes which can break up the web formed by the pectins in the must, as this reduces the viscosity of the liquid and allows the particles to rise up more easily.



1 kg 15 kg

QI UP XC

Fining and stabilizing agent

Qi'Up XC is a formula based on chitin-derived non-animal-origin biopolymers for use in conducting flotation on white, rosé and red wines. It's an all-new flotation additive: natural, biodegradable, non-allergenic and vegan-friendly. Qi'Up XC increases the speed and efficacy with which particles separate from the suspension, regardless of the type of must or the flotation method employed.

Qi'Up XC provides a natural alternative to the use of animal-origin products such as gelatin.

Composition: chitosan from Aspergillus niger and tartaric acid.


Main applications: clarification and stabilization.

• For white and rosé musts: 3-10 g/hL

• For red must after thermovinification: 10-15 g/100 hL

Ion requirements of a chitin derivative compared with two gelatins used for flotation. This measurement is based on the principle of quantifying the charges on the surfaces of the colloids in a specific environment. We can see that the chitin derivatives have a greater charge than the gelatins and therefore display a much greater capacity to agglomerate particles and subsequently form flocs.

This is why Qi'Up is a highly valuable alternative to the use of animal-origin products such as gelatin.



VEGAN-FRIENDLY CHITOSAN-BASED FINING AGENTS





Fining and stabilizing agent for flotation

Qi Up XC Mes is a highly efficient flotation adjuvant in liquid form. Contains a significant quantity of chitosan in a colloidal suspension together with a form of tartaric acid distinguished by its high surface density charge at the pH of the wine; this gives it significant affinity for and reactivity to suspended particles, leading to very swift flocculation. Qi Up XC Mes optimizes thiol production by reducing the concentration of ortho-diphenols, which could form quinones sometimes defined as "aroma traps" under the influence of oxygen.

Composition: chitosan from Aspergillus niger in a colloidal solution and tartaric acid.

Main applications: fining and flotation.

Dosage:

• 20-100 mL/hL

QI UP XC MES

OTHER PRODUCTS

1 kg

10 kg

PK SOL M

Fining and stabilizing agent



Fining and stabilizing agent based on chitin in synergistic co-formulation with PVPP and isinglass. PK Sol M reduces the risk of formation of browning and protein-tannin compounds by rapid flocculation and sedimentation. PK SOL M is also able to reduce the effects caused by poor storage and tone down notes that are too pronounced and/or the presence of bitter flavour, restoring the

aromatic richness of the original product.

Composition: PVPP, chitosan from Aspergillus niger, isinglass.

Main applications: clarification and stabilization of white, red and rosé wines.

Dosage:

• 10-60g/hL depending on the circumstances and operating conditions.







VEGAN-FRIENDLY CHITOSAN-BASED FINING AGENTS



PK SOL M2

Fining and stabilizing agent



(K)



PK SOL M2 is a formulation designed to prevent future oxidation phenomena, colour decay and the formation of bitter tastes. PK SOL M2 is also able to reduce the effects caused by poor storage; by acting on the oxidised components it makes it possible to tone down notes that are too pronounced and/or the presence of bitter flavour, restoring the aromatic richness of the original product.

Composition: PVPP, chitosan from Aspergillus niger, pea protein.

Main applications: clarification and stabilization of white, red and rosé wines.


• 10-60g/hL depending on the circumstances and operating conditions.




The clarifying effects of PkSoIM and PkSoIM2 make it possible to achieve clear wines quickly. Their efficacy comes from the components making up their formulas: the restorative power of chitin derivatives on the colour of oxidised wines combined with a number of compounds providing clarifying effects leads to exceptionally clean, stable wines. PkSoIM2 also contains a top-quality isinglass for even brighter wines.

The PkSoIM2 formulation is suitable for vegan diets.

NTU of a white wine and a rosé wine treated with 20g/hl of PkSoIM and 20g/hl of PkSoIM2



Blind tasting with 8 expert tasters 7 out of the 8 tasters judged PkSoIM and PkSoIM2 to be the best formulations.



VEGAN-FRIENDLY CHITOSAN-BASED FINING AGENTS

10 kg

LUMYCLEAN PLUS

Stabilizer for must and wine





Lumyclean Plus can be used on both must and wine to remove oxidized and oxidizable polyphenols. When making white and red wines, it can be used both as a preventive and a curative treatment in order to prevent oxidative degradation or to 'rejuvenate' oxidized wines, helping them regain their freshness. Lumyclean Plus is also effective at reducing the riboflavin content of white and rosé wines, helping the winemaker avoid the so-called 'light-struck taste' phenomenon. Lumyclean Plus has also been formulated as a defence against the 'light-struck taste' which can occur in white and rosé wines bottled in colourless glass. This defect is the result of the photoreduction of riboflavin (a photosensitive molecule) following exposure to natural or artificial light.

Composition: activated vegetable charcoal, silica gel.

Main applications: removal of oxidized and oxidizable polyphenols, reduction in riboflavin content.

Dosage:

• 5–10 g/hl, depending on needs and working conditions.





FINING AGENTS (PROTEINS)

ALTERNATIVE SOLUTIONS



FYNEO

Yeast-derived fining agent

Fyneo is a protein-rich yeast extract which has undergone a rigorous selection and purification process. It displays exceptional clarifying ability and brings about rapid sedimentation. The proteins extracted from the yeast are highly concentrated and have a high molecular weight (over 15 KDa), which makes them ideal for creating clearer


wines. Fyneo refines the wines by eliminating any hard or bitter notes, while at the same time preserving their positive aromatic profile.


Composition: protein-rich yeast extract.

Main application: clarification of white, red and rosé wines.

- for white and rosé wines: 5-15 g/hL.
- for red wines: 5-30 g/hL.

Fining of a red wine (Merlot) produced using thermovinification (Languedoc Roussillon).





Analysis of turbidity (in NTU) after 14 days of fining at room temperature in the laboratory.

Sensory analysis after 14 days of fining at room temperature in the laboratory (blind tasting performed by 6 expert wine tasters).

FINING AGENTS

PRODUCTS CONTAINING PORCINE GELATIN

20 kg

ATOMIZED GELATIN (COLD SOLUBLE)

Fining agent

This gelatin is particularly suited when, together with fining, the primary purpose is to achieve a good tannin balance. Indeed the action of Atomized Gelatin is particularly effective in the removal of tannins responsible for astringency and a bitter taste. During must clarification, when combined with bentonite or silica sol, we achieve must clarification and an effective reduction of phenolic substances in general.

Composition: porcin gelatin.

Main applications: clarification and stabilization of white and rosé wines lacking in brightness and softness.

Dosage:

• 5-25 g/hL.

1 kg 25 kg

POWDERED GELATIN (HOT SOLUBLE)

Fining agent

In white and rosé wines, treatment with Powdered Gelatin, combined with bentonite or silica sol (xiles), easily removes unstable oxidised phenolic compounds, often the cause of a reduction in chromatic characteristics.

Furthermore, thanks to its high gelling power, it is particularly suitable for the clarification of white and red must.

Composition: porcine gelatin.

Main application: rapid drop in turbidity and formation of compact sedimentation.

Dosage:

- 5-15 g/hL in musts and red wines.
- 3-10 g/hL in white and rosé wines.

ISTANT GEL

Fining agent

The action of Istant Gel is particularly directed to the phenolic substances, in this way the clarification makes it possible to achieve greater stability in regard to oxidative alterations. In wines that are excessively astringent, the use of Istant Gel allows to remove the tannins responsible for this flaw, by bringing more balance to the organoleptic profile of the product.

Composition: 30% porcine gelatin in an aqueous solution, SO₂.

Main application: suitable for clarifying treatments of musts and wines.

Dosage:

- 5-15 g/hL as an adjuvante for clarification.
- 15-30 g/hL as a colour stabilizer in red wines.
- 30–80 g/hL as tannin-remover according to the content of tannin.









25 kg







ISTANT GEL 45



25 kg

Fining agent

Istant Gel 45 carries out its action by reacting with tannins present in musts and wines. The interaction causes the formation of large floccules that fall, allowing the clarification of the product. Instant Gel 45 is therefore a valuable adjunct to be used in the process of must clarification, especially if it is particularly tannic, or to complete a spontaneously insufficient clarification. Its use has proved valuable in situations where normal gelatins are unsatisfactory.

Composition: 45% porcine gelatin in an aqueous solution, SO₂.

Main application: suitable for clarifying treatments of musts and wines.

Dosage:

• 5-30 g/hL.

PRODUCTS CONTAINING ISINGLASS

1 L 10 L



CRISTALLINE LIQUID

Fining agent containing isinglass

Adding Cristalline to wine enables the flocculation and sedimentation of suspended particles. Together with a long decantation process, it constitutes one of the best methods for fining and stabilizing wines. It also increases filterability.





1 kg

Composition: isinglass.

Main applications: clarification and stabilization of white and rosé wines.

Dosage:

• 10-12 cL/hL.

CRISTALLINE PLUS

Fining agent containing isinglass

Adding Cristalline Plus to wine enables the flocculation and sedimentation of suspended particles. Together with a long decantation process, it constitutes one of the best methods for fining and stabilizing wines. It also increases filterability.



Main applications: clarification and stabilization of white and rosé wines lacking in brightness and softness.

Dosage:


• 1,5-3 g/hL. Before fining with isinglass, it may be necessary to treat the wine with Tanin TC.


PRODUCTS CONTAINING EGG ALBUMIN

1 kg 25 kg

OVOCLAR

Fining agent made from egg albumin





Main applications: organic fining agent for treating wines.

Dosage:


- 5-10 g/hL in white wines.
- 10-25 g/hL in red wines.

COLLOIDAL SILICA

25 kg - 1300 kg in bulk

XILES 40

Colloidal silica



Composition: 40% colloidal silica dioxide solution.

Main application: used in fining processes on white must and wine.

Dosage:

- 30-70 g/hL in must, together with 15-30g/hL of Istant Gel.
- 15–30 g/hL in wine, together with 3–10g/hL of Istant Gel.
















Winemakers are well aware of the need to stabilize tartrates, colloids, proteins and colour. Perdomini-IOC provides a complete range of stabilization aids for each contingency that can arise in a certain wine and/or when using a certain stabilization method

Unstable colloids

GUM ARABIC: STABILIZATION/PROTECTION

Gum arabic and colloid stabilization

When the wine is ready for bottling, the winemaker needs to perform chemical, sensory and stability tests in order to be sure that the wine will maintain its level of quality over time. Gum Arabic is an important ally in the quest to stabilize wines. It's a natural exudate from the bark of the acacia tree and is used as a 'stabilizer' for unstable colloids in wines. It improves their overall aromatic profile and mouthfeel, intensifies sensations of sweetness and roundness in the mouth and reduces astringency. It also reduces the risk of iron and copper casse marring the wine.



GUM ARABIC: STABILIZATION/PROTECTION

As it is a protective colloid, it combats the precipitation of suspended particles and absolutely must be used on "pre-setting" wines, i.e. wines which have been clarified and stabilized or are at the final stages of filtration. The gum arabic should be incorporated into the wine the day before the final filtration before bottling, or after the final filtration, in this case using a high-precision metering pump. Regulation EU 2019/934 lays down that the maximum dosage of gum arabic allowed is 30g/hl of dry matter. The precise dosage varies according to the desired sensory results and the level of instability in colouring substances measured after fining and initial filtration. To measure this instability, we recommend performing a cold-resistance test by chilling a sample of the wine to 4°C for 48 hours.

Colouring substance stability test

- Measure the turbidity of the initial sample (NTU before exposure to cold).
- If the turbidity is > 2 NTU, filter 30ml using a 0.65µm membrane.
- Put these 30ml (whether filtered or not) into a bottle and leave at 4°C for 48 hours.
- At the end of this time, shake the bottle then, after 15 minutes at room temperature, measure the turbidity of its contents (NTU after exposure to cold).

Δ NTU = NTU after cold - NTU before cold

Adding gum arabic is highly useful in wines with instability levels under $30\,\Delta\,\text{NTU}.$



Stabilizat

STABILIZATION


GUM ARABIC: STABILIZATION/PROTECTION

10 kg - 25kg 200 kg - 1000 kg



READY GUM ELITE

Stabilizing gum arabic



Ready Gum Elite helps avoid precipitation of colour particles and lend wines greater drinkability, roundness and smoothness.



Main application: for treating not only white, red and rosé wines but also fortified wines, in any situation where the winemaker wants to avoid precipitation of colour particles after bottling.

Dosage:

- 30-80 g/hL to avoid precipitation of colour particles.
- 50–100 g/hL to achieve the desired sensory characteristics.

Ready Gum Elite:

the levorotatory gum arabic that can be used in microfiltration.




TABLE OF SUMMARY OF THE MAIN SENSORY CHARACTERISTICS AND APPLICATIONS

7	ReadyGum 20	ReadyGum ELITE	ReadyGum L
Softness		0000	0000
Body	000	0000	0000
Tartrate stability	0000	O	00
Colour stability	O	0000	0000

STABILIZATION GUM ARABIC: STABILIZATION/PROT

10 kg - 25 kg 200 kg - 1000 kg



READY GUM L

Stabilizing gum arabic



Ready Gum L acts by coating the hydrophobic micelles, thereby preventing them from binding together and forming colloids big enough to potentially cause physical instability problems. Ready gum L helps protect wines against ferric-phosphate and oxydase casse.



Composition: gum arabic from Acacia Kordofan in 20% water solution, SO₂.

Main application: for treating not only white, red and rosé wines but also fortified wines, in any situation where the winemaker wants to avoid precipitation of colour particles after bottling.


Dosage:

- 30-60 g/hL to avoid precipitation of colour particles.
- 50–100 g/hL to achieve the desired sensory characteristics.

1 kg

DRY GUM L

Levorotatory gum arabic in powder



The unique polysaccharide nature of Dry Gum L means it lends greater structure and more rounded aromas and flavours to the wine with very little clogging.

Composition: gum arabic from Acacia Kordofan.

Main application: protective colloid against colour instability recommended for treating red, rosé and fortified wines, in any situation where the winemaker wants to avoid precipitation of colour particles after bottling.

Dosage:

• 10-100 g/hL.














GUM ARABIC: TARTARIC STABILIZATION


10 kg - 25 kg 200 kg - 1100 kg



READY GUM 20

Stabilizing gum arabic





Composition: gum arabic from Acacia Seyal in 20% water solution, SO₂.

Main application: recommended for treating white, red and fortified wines, whenever it is necessary to avoid clouding, flocculation or precipitation after bottling.

Dosage:

- 40–100 g/hL to prevent tartrate instability.
- 50-200 g/hL or more to achieve the desired sensory characteristics.



Dextrorotatory gum arabic in powder

Dry Gum R acts as a protective colloid by wrapping around hydrophobic micelles and stopping them from aggregating into macromolecules large enough to cause physical instability in the wine. Its formulation as a powder means that the winemaker can enjoy all the advantages of a high-quality gum arabic without the addition of water, SO₂ or citric acid.

Composition: gum arabic from Acacia Seyal.

Main application: recommended for treating white, red, rosé and fortified wines, whenever it is necessary to avoid clouding, flocculation or precipitation after bottling.

Dosage:

• 10-100 g/hL.







1 kg 10 kg





TARTRATE AND CALCIUM STABILIZATION

1 kg 15 kg

METACREMOR 40+

Metatartaric acid

Tartrate stabilizer that acts by wrapping around tartrate crystals, thereby stopping the growth of microcrystals and their subsequent precipitation. Its high degree of esterification guarantees great efficacy on both potassium bitartrate and calcium tartrate, so that the wine will remain stable for the whole winter period.





Composition: metatartaric acid.

Main application: tartaric stabilization.

Dosage:

• 10 g/hL max.

1 kg 15 kg

METACREMOR INSTANT

Metatartaric acid

Tartrate stabilizer formulated in response to the needs of wineries where it's impossible to wait 48 hours or more before bottling. Metacremor Instant avoids the problem of clouding in wines after metatartaric acid with a high degree of esterification (> 40) has been added (although this is a totally normal phenomenon that disappears within 48 hours).



Composition: metatartaric acid.

Main application: tartaric stabilization.

Dosage:

• 10 g/hL max.

CREME DE TARTRE

Crystallizer

Composition: micronized potassium bitartrate.

Main application: induce bitartrate crystal precipitation when cold-stabilizing wines.

Dosage: varies according to needs and the tartrate-stabilization system employed.



TARTRATE AND CALCIUM STABILIZATION

5 kg

TARTRATE DE CALCIUM

Tartaric stabilizer

Calcium tartrate makes it possible to achieve effective tartar stabilization by stabilizing both potassium bitartrate and calcium tartrate in a single step.

Composition: purified calcium tartrate tetrahydrate.

Application: encourage tartar precipitation and help stabilize the tartrates in the wine by decreasing the final concentration of potassium bitartrate and calcium tartrate.

Dosage:

• 100-200 g/hL

DUOSTAB

Tartaric stabilizer

Duostab has been specially designed for use in cases where tartar stabilization is tricky and both forms of tartrate salt - potassium bitartrate and calcium tartrate - are involved. Duostab is a technological additive containing potassium bitartrate and calcium tartrate in optimal proportions for allowing cold stabilization of calcium tartrates in a single step.

Composition: potassium bitartrate and calcium tartrate.

Application: encourage tartar precipitation and help stabilize the tartrates in the wine by decreasing the final concentration of potassium bitartrate and calcium tartrate.

Dosage:

• 100 g/hL - 400 g/hL

CRISTAL CREMOR S

Tartaric stabilizer

Using Cristal Cremor S is an effective way to optimize the practice of stabilizing wines through refrigeration, as it ensures tartrate stability and speedy completion.

Composition: combined potassium tartrates to boost the effects of superfine diatomaceous earth.

Main application: adjuvant for cold stabilization of tartrates.

Dosage:

• 30-60 g/hL depending on the total acidity of the wine to be treated.

15 kg

25 kg

1 kg

CARBOXYMETHYLCELLULOSE

STABILIZATION

CRYSTAL BALANCE

Carboxymethylcellulose (CMC)

Crystal Balance is a protective colloid that inhibits the formation (or nucleation) and subsequent growth of tartrate crystals in wine. It's a polysaccharide with a delicate balance between its degree of polymerization, degree of substitution and uniformity, so that the wine will be successfully stabilized with a low viscosity and very little gel formation.

Composition: carboxymethylcellulose.

Main application: adjuvant for tartrate stabilization.

Dosage: the dosage generally used ranges from 5-10g/hL, with a maximum permitted dosage of 10g/hL.

20 kg

CRYSTAL BALANCE SOL. 5%

Carboxymethylcellulose (CMC) in solution

Crystal Balance sol. 5% is a protective colloid that inhibits the formation (or nucleation) and subsequent growth of tartrate crystals in wine. Crystal Balance sol. 5% comes in a readymade solution for speed and ease of use. Not recommended when the wine has been or will be treated with lysozyme.


Composition: carboxymethylcellulose, SO₀.

Main application: adjuvant for tartrate stabilization.

Dosage: 100-200 mL/hL

Maximum permitted dosage: 200 mg/L.

One of the main properties of CMC is that it can inhibit the formation of tartrate microcrystals. Acting as a protective colloid, it blocks the growth of crystals and stops tartrate salts from precipitating. It's not advisable to add CMC to red wines, as it interacts with the phenolic compounds. When used on white wines, the wine must display total protein stability. We recommend performing a preliminary heat test to be sure that the proteins in the wine are stable. CMC is incompatible with wines treated with lysozyme. It's advisable to add the CMC to the wine 48 hours before the final filtration before bottling, otherwise it could clog up the filter cartridges. The maximum dose allowed by law is 100mg/L.



TREATMENTS TO CORRECT SPECIFIC DFFFCTS

DECOLOURIZING CARBON

CARBO GRANÈ

Decolourizing carbon in pellet

Carbo Grané is a carbon in pellet form with an exceptional decolourizing ability. The chemical activation process leads to extremely porous particles which can adsorb various different types of molecules, depending on their size. The careful selection of ingredients is a guarantee of the quality of Carbo Grané, and of a total absence of unwanted by-products, such as metals or odours foreign to the treated product.

Composition: activated vegetable carbon.

Application: Carbo Grané is particularly effective at decolourizing wines.

Dosage:

• 100 g/hL - 400 g/hL

CARBONE DECOLORANTE SUPERATTIVO CP

Vegetable decolourizing carbon

Activated vegetable carbon boasting highly porous particles which can adsorb various different types of molecules, depending on their size. During fermentation: to improve colour and stability in the wine. When added to white musts, Carbone Superattivo CP eliminates any excess anthocyanins, leading to improvements in the colour profile of the wine. When used during fermentation, it helps remove any pesticide residues, leading to a smoother fermentation process.

Composition: activated vegetable carbon.

Main application: decolourization.


Dosage:

• 10-100 g/hL.


TREATMENTS TO CORRECT SPECIFIC DEFECTS **DECOLOURIZING CARBON**

NO DUST CARBON

Dust-free decolourizing carbon







Main application: decolourization.

Dosage:

• 10-100 g/hL.

10 kg

TOPCARBON

Vegetable decolourizing carbon

The adsorption capacity of Top Carbon makes it highly effective on white wines marked by excessively intense colouring, which may be caused by oxidation or maderization phenomena. Top Carbon can also be used successfully to reduce the iron content in wines, helping protect them against iron casse.

Composition: vegetable activated carbon.

Main application: decolourization.

Dosage:

• 10-100 g/hL.

1 kg 10 kg

CARBO F

1 kg

Vegetable decolourizing carbon

Special vegetable carbon with intense decolourizing power. During production, Carbo F undergoes a chemical-physical activation process that increases its efficacy, as it gives the carbon extra reactivity. Carbo F is highly porous, making it perform extremely well in winemaking, as the contact surface between the carbon and the must/wine is exceptionally large. The low doses required allow the wine to preserve its whole bouquet of aromas.



Main applications: decolourize white wines marred by excessively intense colour and reduce iron content. Can also be used to decolourize red musts.

Composition: vegetable carbon.

Dosage:

• 10-100 g/hL










DECONTAMINATING CARBON

10 kg

EVF - FREE

Off-odour adsorbent





Composition: active carbon and silica gel.

Main application: eliminate off-odours, in particular musty and/or phenolic odours.

Dosage:

• 30-100 g/hL

LUMYCLEAN REX





1 kg



Lumyclean Rex is a blend of a special activated vegetable charcoal - selected for its high level of purity – and silica gel. Effective at very low doses and with short contact times, Lumyclean Rex is a special treatment for white and rosé wines that helps winemakers avoid the so-called 'lightstruck taste' caused by the photoreduction of riboflavin (a photosensitive molecule) following exposure to natural or artificial light when the wine has been bottled in colourless glass. If the correct dosages are used, Lumyclean Rex does not alter the colour of rosé wines.

Composition: activated vegetable charcoal, silica gel.

Main application: reduction in riboflavin content.

Treatment to combat 'light-struck' defects

• 3–5 g/hl, depending on needs and working conditions

TREATMENTS TO CORRECT SPECIFIC DEFECTS

ANTI-REDUCTION TREATMENTS

1 L 5 L

REDOXYL

Treatment for reduction faults



Redoxyl eliminates the unpleasant odours and sensations of reduction caused by sulphurous compounds (SH).



Composition: copper sulphate, citric acid, potassium metabisulphite.

Main application: eliminate sensations of reduction.

Dosage:

• 8-10 ml/hL

Legal limit: 10 mL/hL, as long as the wine treated does not have a copper value more than 1 mg/L.

SOLUTION 700





This product targets sensations of reduction in still wines, bases for sparkling wines and sparkling wines at the time of the tirage.



Composition: copper sulphate in solution, citric acide and SO₂.

Main applications: prevention and treatment of sensations of reduction.

Dosage:


- Still wines: 2-4 cL/hL.
- Bases for sparkling wines: 1-2 cL/hL.
- Sparkling wines: 1–2 cL in the quantity of liqueur d'expedition necessary for 100 bottles.

ACIDIFIERS

1 kg 25 kg

ACID CITRIQUE

Acidifier



Dosage: by law, the amount in the finished wine must not exceed 1g/L.



TREATMENTS TO CORRECT SPECIFIC DEFECTS ACIDIFIERS

ACIDE LACTIQUE

Acidifier

Acide Lactique is of natural origin, obtained by sugar lactic fermentation.

Composition: L-lactic acid 88% in aqueous solution.

Main applications: it is used to acidify musts derived from extremely ripe and insufficiently acidic grapes, or in wines.

Dosage:

- In white and rosé wines: from 3 to 15 g/hl for freshness and fineness.
- In red wines: 10-20g/hl to boost freshness in the finish and reduce all forms of bitterness

Maximum legal dosage and acidification declaration according to current regulations.

ACIDE MALIQUE D,L

Acidifier

DL-malic acid makes it possible to increase and manage the total acidity of wines.

Composition: DL-malic acid.

Main application: useful to acidify juices extracted from very ripe, insufficiently acidic grapes.

Dosage: maximum dose allowed under the laws of the region in question.

ACIDE TARTRIQUE L+

Acidifier

Composition: L(+) tartaric acid.

Dosage: maximum dose allowed by the law in force.

1 kg - 5 kg 25 kg



5 kg 25 kg

5 kg

25 kg



TREATMENTS TO CORRECT SPECIFIC DEFECTS

DE-ACIDIFIERS

1 kg 25 kg

CALCIO CARBONATO

De-acidifier

Calcium carbonate is a deacidifier suitable for use in must and wine. It induces the precipitation of tartaric acid in the form of an insoluble salt.

1 kg

25 kg

Main applications: deacidify must and wine.

Composition: calcium carbonate.

Dosage:

• 50 g / hL of calcium carbonate reduces acidity by 0.5 g/L, expressed as sulphuric acid.

BICARBONATE DE POTASSIUM

De-acidifier

Main applications: deacidify must and wine.

Composition: potassium bicarbonate.

Dosage: under normal conditions, adding 1 g/L of potassium bicarbonate reduces acidity by 0,49 g/L, expressed as sulphuric acid.

1 kg 25 kg

TETRACOMPLEX

Composite deacidifier

The components making up Tetracomplex show a tendency to react with the tartaric acid in the wine, neutralizing its acid groups and transforming it into its insoluble salts, potassium bitartrate and calcium tartrate. In this way, Tetracomplex allows the winemaker to reduce any excess acidity and attain a marked improvement in the sensory balance of the treated wine. Moreover, reducing acidity – especially in red wines – provides a significant boost to the start-up of malolactic fermentation.

Composition: potassium tartrate, potassium bicarbonate, calcium carbonate.

Main applications: wines with excessive total acidity and abnormally low pH.

Dosage: according to needs, calculating that 100g/hL of product reduces the total acidity by approximately 1‰.

02-





TREATMENTS TO CORRECT SPECIFIC DEFECTS

OTHER ACIDS

1 kg 25 kg

ACID ASCORBIQUE

Antioxidant preservative

Composition: L-ascorbic acid.

Main applications: prevent wine oxidation.

Dosage: legal limit: 25 g/hL





MICROBIOLOGICAL STABILIZATION

SULPHITES

OXYLESS RANGE

1 kg

OXYLESS U

Special antioxidant for grapes

A combination of antioxidant products that prevents oxidation thanks to its components. In particular, L-ascorbic acid eliminates any oxygen present and, at the same time, limits the action of oxidase enzymes such as laccase and polyphenol oxidase. The simultaneous presence of gallic tannins and sulphur dioxide makes it possible to block the action of oxygenated water and any free radicals. The presence of proanthocyanidin tannins, rich in flavonoids, makes it possible to obtain fruitier and more complex wines.

Composition: potassium metabisulfite, L-ascorbic acid, gallic tannin and proanthocyanidin tannin rich in flavonoids.

Main application: prevents grape oxidation during harvesting, unloading and pressing.

Dosage:

- 10-25 g/hL for white grapes
- 20-30 g/hL for red grapes.

500 g 10 kg

OXYLESS M

Special must antioxidant

A combination of antioxidant products that, thanks to its components, makes it possible to obtain fresh fruity wines with greater longevity. The presence of yeast cell walls with quick release of glutathione increases the shelf life of the wine. The absence of SO₂ and the simultaneous presence of gallic tannins and proanthocyanidins in the formula of OxyLess M enable the glutathione to be maintained in non-oxidized form. The use of OxyLess M at the end of alcoholic fermentation enables glutathione to be preserved in the must. The simultaneous presence of tannin contributes to an improvement in the structure and taste balance of the final wine.

Composition: yeast cell walls rich in glutathione (GSH) and sulphurated amino acids, gallic tannin and proanthocyanidin tannin rich in flavonoids.

Main application: protects varietal and fermentative aromas and flavours in white and rosé wines.

Dosage:

- 20-30 g/hL in white grape must
- 10-20 g/hL in red grape must.









25 kg

Preservative - Antioxidant

Main application: provide microbiological and antioxidant protection.

Composition: potassium metabisulfite.

• 1 g of potassium metabisulfite releases 0.5g of SO₂. Maximum dosage usually apply, please refer to the applicable legislation.

48 capsules

INODOSE 2



• one capsule releases 2 g of SO₂.

42 capsules

INODOSE 5

Capsules containing potassium metabisulfite that release 5 g of SO₂ per capsule. Designed to simplify sulphuring operations.

Composition: potassium metabisulfite.

Application: correct the quantity of SO₂ in the must.

Dosage:

STERISOLFITO 15

Sulphite in liquid form

As it's a sulphite, Sterisolfito 15 performs a preventive task against the dangerous oxidation reactions that can take place in must and wine. It allows the winemaker to combat both oxidation problems and the growth of indigenous microflora.

100g of product releases around 15g of SO₂).

Main application: prevents oxidation.

Dosage:

• 60-80 g/hL in must extracted from compromised grapes.

10 kg

OXYLESS V

Special antioxidant for wine

A combination of antioxidant products that prevents oxidation during racking operations. The presence of L-ascorbic acid eliminates any oxygen present and makes it possible to reduce the amount of SO₂ needed to protect against oxidation. The simultaneous presence of sulphur dioxide makes it possible to block the action of oxygenated water and any free radicals. The green tea proanthocyanidin tannin, rich in flavonoids, makes it possible to improve the structure and body of the wine, stabilize the colouring matter and speed up the consumption of dissolved oxygen during racking.

Composition: potassium metabisulfite, L-ascorbic acid, green tea proanthocyanidin tannin rich in flavonoids.

Main application: protects varietal and fermentative aromas and flavours in white and rosé

Dosage:

• 10-20 g/hL.

Before each racking, add OxyLess V to stop too much oxygen getting into the wine.

OTHER SULPHITES

FERROBLOCK

Redox-reaction stabilizer

By lowering the redox potential of the wine, Ferroblock prevents the onset of oxidation reactions that can make the wine lose freshness and liveliness. Excellent results have been achieved in preventing and treating colour loss caused by oxidase casse. When Ferroblock is added to the wine before its last filtration before bottling, it ensures long-lasting stability and plays a decisive role in creating fresh, delicate, intensely fruity wines.

Composition: anhydrous citric acid, potassium metabisulfite, L-ascorbic acid.

Main application: prevents wine oxidation.

Dosage:

- 10-25 g/hL for white wines.
- 10-20 g/hL for red and rosé wines.

Adding 10g/hL increases the total SO₂ level by approximately 20mg/L.


















Sulphite

Capsules containing potassium metabisulfite that release 2g of SO₂ per capsule. Designed to simplify sulphuring operations.

Composition: potassium metabisulfite.

Main application: correct the quantity of SO₂ in the must.

Dosage:

Sulphite

• one capsule releases 5 g of SO₂.

Composition: ammonium bisulfite, enough deionized water for a final volume of 100 ml (every

- 30-40 g/hL in must extracted from healthy grapes.

MICROBIOLOGICAL STABILIZATION SULPHITES

STERISOLFITO 45

Sulphite

Sterisolfito 45 inhibits the multiplication of indigenous blastomycetic and bacterial flora, thereby ensuring that only the selected starters initiate fermentation and bring it to completion. Sulphuring with ammonium bisulfite has an advantage over using potassium metabisulfite in that it shifts the pH towards slightly lower values. This property can be highly valuable in winemaking situations where the acidity of the wine needs to be preserved as far as possible.

Composition: ammonium bisulfite, enough deionized water for a final volume of 100 ml (every 100g of product releases around 45g of SO_2).

Main application: prevents oxidation.

Dosage:

- 10-15 g/hL in must extracted from healthy grapes.
- 20–25 g/hL in must extracted from compromised grapes.

SULFITANIN

Sulphite

Ammonium bisulfite in solution with a concentration of 100g/I of SO_2 . Also contains tannin, which boosts the antiseptic and antioxidant properties of the sulphur.

Composition: ammonium bisulfite, enough deionized water for a final volume of 100 ml (every 100g of product releases around 45g of SO₂).

Application: sulphur additive for must.

Dosage:

• 5-10 cL/hL

25 kg





1 L

11/11



MICROBIOLOGICAL STABILIZATION

OTHER SULPHITES

Penny Damigiane 480 tablets
Penny Fusti 120 tablets
Penny Vasche 50 tablets

PENNY

Floating anti-Flowers-of-Wine tablets



The air enclosed between the surface of the liquid and the lids of wine vessels is the perfect breeding ground for the mycetic microflora that leads to Flowers of Wine. This growth inevitably spoils the wine, and if left unchecked can lead to sourness and even acescence. Allyl isothiocyanate displays an excellent capacity to inhibit the development of oxidizing yeasts. Penny tablets are available in three different formats:

- Penny Damigiane: for relatively small vessels, but with a capacity of over 20 litres;
- Penny Fusti: for medium-sized vessels;
- Penny Vasche: for large vessels.

Composition: solid paraffin and allyl isothiocyanate.

Main application: prevent the onset of Flowers of Wine and the development of oxidizing yeasts.

Dosage:

- Penny Damigiane: 1 tablet, to be replaced every 2-3 weeks.
- Penny Fusti: 1-2 tablets, to be replaced every 15-20 days.
- Penny Vasche: 1-2 tablets, to be replaced every 15-20 days.

1 kg 25 kg

SORBENOL

Fungistatic stabilizer



The anti-fermentation action of Sorbenol is particularly marked when it comes to Saccharomyces cerevisiae, which is why it's advisable to use it in cases where the winemaker wants to reduce the risk of refermentation to a minimum – for example, in wines with sugar residues or in partially fermented musts. Sorbenol is also effective at inhibiting the development of the microorganisms responsible for Flowers of Wine.

Composition: potassium sorbate.

Main Application: avoid refermentation and the onset of Flowers of Wine.

Dosage: 27g/hl of Sorbenol equals 200mg/l of sorbic acid in the wine.



TANNINS


Exogenous winemaking tannins are polyphenols. Extracted from naturally occurring vegetable sources, they come from various botanical species such as gall, wood (chestnut, oak, mimosa, exotic woods and quebracho) and grapes (grape-pip and skin tannins).

Ellagic tannins ("Ellagitannins") and gallic tannins ("Gallotannins") belong to the category of hydrolyzable tannins: when exposed to an acid or to heat, they lead to the formation of ellagic or gallic acid.

Proanthocyanidin tannins belong to the category of condensed tannins.

Depending on their source – which determines their properties as winemaking aids –, they can have various uses:

- as fining agents, because the tannins interact with the proteins in the wine and the resulting compound precipitates;
- as antioxidant and antioxydase agents (as they consume O_a);
- as colour stabilizers (they limit the oxidative degradation of anthocyanins and facilitate co-pigmentation)
- to help improve the structure of the wine:
- to eliminate the reduction smell.



AGING PRE-BOTTLING

Approximate dose: from 10 to 40 g/hL

CHARACTERISTICS OF THE WINE

Approximate dose: from 1 to 5 g/hL **TANNINS**


LIMITING ANTIOXIDANT AND ANTIOXYDASE EFFECTS

1 kg 5 kg



ESSENTIAL ANTIOXIDANT

Antioxidant gallic tannin





Composition: gall-nut tannin.

• 10 g/hL of must or wine.

Main application: inhibit the enzymatic activity responsible for oxidation in must.

Tannin selected as part of the project







0,5 kg 5 kg

GALLOTAN

Dosage:

Gallic tannin for fining and balanced mouthfeel

Thanks to its strong reactivity to proteins, Gallotan is able to avoid the problem of overfining when the wine is clarified using protein-based adjuvants (e.g. gelatin). Gallotan does not alter the wine's colour and is recommended for use above all on white wines which, as they naturally contain less tannins, are more susceptible to protein instability.





Composition: gall-nut tannin.

Main application: inhibit the enzymatic activity responsible for oxidation in must.

Dosage:

• 3-15 g/hL.

Perform laboratory tests in order to establish the optimal dosage.

Thanks to its intense antioxidant effect, Tan FlavourFF protects both the colouring substances and the aromatic components from oxidation so that the wine does not suffer from browning. Moreover, using Tan Flavour FF in the early stages of winemaking allows the addition of specific



Composition: granular proanthocyanidin tannin derived from lemon and acacia wood.

aroma precursors that intensify the aromas of flowers and fruits in the wine.

Main application: protects colouring substances and aroma producers at the same time as providing fruity and floral aroma precursors.

- white wines: 4-10 g/hL, divided into two doses, the first on the must and the second 24 hrs after initiating fermentation.
- red wines: 10–15 g/hL in a single dose 72 hrs after initiating alcoholic fermentation.

IMITING THE OXIDATIVE DEGRADATION OF ANTHOCYANINS; PROTECTING AND STABILIZING COLOUR OVER TIME

1 kg 5 kg

CROMOX G

Tannin providing antioxidant and colour-stabilizing power

Cromox G protects the colouring matternt substances that has just been extracted from the grapes. In synergy with SO₂, Cromox G acts to protect colouring substances from the outset, an action which can continue during maceration through Cromofix.

Cromox G can also be used when aging red wines, as it speeds up the evolution of the polyphenols in the wine (especially when pre-used casks are being used), and can prevent the reduction smell. Lastly, its high level of reactivity to proteins makes Cromox G a valuable tool for avoiding overfining.

Composition: blend of hydrolyzable and condensed tannins.

Main application: colour stabilization in red wines during pressing and in the earliest stages of maceration.

Dosage:


• 10-30 g/hL.

TANNINS LIMITING, PROTECTING AND STABILIZING

5 kg

CROMOFIX SR

Tannin for colour stabilization



Composition: condensed tannin.

Main applications: colour stabilization in red wines during maceration and winemaking.

on Botrytis-infected grapes, as it has the capacity to inactivate laccase.

Dosage:

- 10-30 g/hL for colour stabilization.
- 15-50 g/hL to inhibit laccase.

1 kg 25 kg

TANIN TC

Tannin for fining

Tanin TC is a hydrolyzable tannic acid extracted from chestnut wood by dissolving it in water. When it reacts with the fining agent, Tanin TC forms a precipitate which, as it settles, drags down the particles clouding the wine along with it. The result is a much clearer wine and one with a greatly improved polyphenol structure.

Composition: tannin extracted from chestnut wood.

Main application: suitable for fining wines in tandem with Cristalline (isinglass) or other protein-based fining agents (pork or fish gelatin).

Dosage: perform laboratory tests before each collage in order to establish the optimal dosage. In general, doses should be around:

• 7-8 g/hL, together with a fining agent.

1 kg - 5 kg 15 kg

TANIN SR

Tannin for colour stabilization

Tanin SR is a catechinic and ellagic tannin extract which presents in the form of a dark-brown powder and is slightly soluble in water. When used before maceration in red-wine production, it stabilizes colour. Adds intensity and structure to finished red wines. At low doses, it can also increase softness in white wines.

Composition: Quebracho-wood tannin.

Main application: stabilize colour during maceration when crafting red wines.

Dosage:

- during maceration: 15-30g per 100kg.
- during winemaking: 15-30g per hL.













TANNINS

LIMITING, PROTECTING AND STABILIZING

FILL COLOR

FULLCOLOR

Blend of tannins and yeast polysaccharides



Fullcolor is a preparation made from proanthocyanidin and ellagic tannins with a high polysaccharide content (from yeast), designed for red wines. It's a 100% natural product that can be used at the fermentation stage to increase phenol

compound stability and reduce tannicity while boosting the body and structure of the wines. The proanthocyanidin tannins found in Fullcolor are similar to the condensed tannins found in grapes; they aid colour stabilization and help avoid colour loss. Ellagic tannins, on the other hand, help protect the anthocyanins, meaning that they are freed from the oxidation and precipitation problems that can arise at the aging stage.

Composition: proanthocyanidin and ellagic tannins and yeast polysaccharides.

Main application: stabilize colour when crafting red wines without causing precipitation.

- for colour stabilization: 20-40 g/hL.
- for laccase inhibition: 40-80 g/hL.
- for a considerable flavour boost: 30-60 g/hL.

VOLUTAN

Grape tannin for fining, colour stabilization and structure

Volutan is a stable, 100% soluble, tannin in liquid form which does not lead to losses through precipitation. When used on red wines, it stabilizes their colour and improves their phenolic potential in terms of both quantity and quality. In rosé wines, Volutan removes excess proteins (boosting the action of bentonite clays), maintains colour and improves resistance to oxidation. Volutan is a 100% natural product obtained exclusively from white wine grapes, with no colourants, no organic solvents and no export restrictions. It displays identical properties to the tannins naturally found in must and wine.

Composition: grape tannin in liquid form.

Main applications: improve clarification, colour stabilization and phenolic potential (structuring effect).

Dosage:

- red winemaking: 15-40mL/hL.
- rosé winemaking: 10-20mL/hL.

TANIFASE ELEVAGE

Oak-wood tannin for colour and aroma stabilization

Taninfase Elevage is a hydrolyzable oak-wood tannin (Quercus robur e petraea) which boosts the preservation and stabilization of the colour and bouquet of wines at the same time as improving their overall balance.

Composition: oak-wood tannin (Quercus robur e petraea).

Main application: colour and aroma stabilization.

Dosage:

• 5g-15g/hL

1 kg 10 kg















1 kg







MANN BOUQUET B19

Optimising alcoholic fermentation of white and rosé musts

Mann Bouquet B19 lends notes of flowers and white fruits at the same time as providing an antioxidant effect leading to greater fullness and sweetness in the mouth.

There are many benefits that can be gained by using this product: enhancement of white fruit (e.g. apple, pear) and floral notes, as well as

increasing softness and volume, balance and structure in the mouth. For even greater expression of aromas and flavours, we recommend using this product in tandem with IOC Révélation Thiols for white must or whith IOC Fresh Rosé for rosé must.

Composition: deactivated yeast rich in glutathione with a high content of soluble mannoproteins, gallic and acacia tannins.

Main application: production of white and rosé wines enhanced whith fruit and floral notes.

Dosage:

• red and rosé musts: 10-40 g/hL.

Maximum dose allowed by the EU: 50 g/hL.

SPECIAL TANNIN-BASED PRODUCTS FOR VINIFICATION

BOUQUET RANGE



MANN BOUQUET R16

Optimising alcoholic fermentation of red and rosé musts





Mann Bouquet R16 protects colour and facilitates anthocyanin-tannin condensation while red-wine grapes are macerating. There are many benefits that can be gained by using this product; protection against oxidation and the effects of oxidase enzymes, stabilization of colouring substances, enhancement of dark berry (especially blackcurrant) and spicy

notes, and greater persistence of fresh and fruity notes over time.

Composition: yeast hulls with a high content of soluble mannoproteins, ellagic and grape seed tannins

Main application: used to produce red and rosé wines with strong spicy and dark-berry notes.

Dosage:

IHANCE THE SPIC

• red and rosé musts: 10-40 g/hL.

Maximum dose allowed by the EU: 80 g/hL.

1 kg







SPECIAL TANNIN-BASED PRODUCTS FOR VINIFICATION

TANIN BO

TANIN BOUQUET B45



1 kg

Tannins for white and rosé musts



The special process of extraction and drying at a low temperature used during the production process of Tanin Bouquet R45 makes it possible to extract from the wood glycosylated forms of aromatic compounds such as terpenes (citronellol, linalool, etc...) and norisoprenoids (beta-dam-ascenone, etc...), responsible for the fruit and floral notes in wine. The

resulting wine therefore has intense aromas of lemon, grapefruit, apple and white flowers, which complement the varietal aromas and those produced during fermentation. The aromatic expression is heightened with the use of certain strains of yeast such as IOC Révélation Thiols (for white musts) and IOC Fresh Rosé (for rosé musts).

Composition: blend of condensed and gallotannins.

Main application: produce white and rosé wines with a real citrussy feel.

Dosage:

• rosé must: 5-15 g/hL.

• white must: 2-15 g/hL.



TANIN BOUQUET R36



1 kg

Tannins for red and rosé musts

The special process of extraction and drying at a low temperature used during the production process of Tanin Bouquet R36 makes it possible to extract from the wood glycosylated forms of aromatic compounds such as terpenes (citronellol, linalool, etc...) and norisoprenoids (beta-dama-scenone, etc...), responsible for the hints of red berry fruit in wine. The

resulting wine therefore has intense aromas of cherry, strawberry, marasca cherry, blueberry, etc. which complement the varietal aromas and those produced during fermentation. The aromatic expression is heightened with the use of certain strains of yeast such as IOC R9008 (for red musts) and IOC Fresh Rosé (for rosé musts). Tanin Bouquet R36 also promotes colour stabilization and prevents oxidation of the primary notes.

Composition: blend of condensed tannins extracted from wood of red berry fruit species.

Main applications: production of red and rosé wines for a red berry fruity taste, with a strong action in stabilising the colour.

Dosage:

- rosé must: 2-15 g/hL.
- red must: 5-20 g/hL.


SPECIAL TANNIN-BASED PRODUCTS FOR VINIFICATION

1 kg

TANNIN BOUQUET B49



A blend of gallic tannins and deactivated yeasts to increase the aroma of white of rosé wines



ing fermentation. The level of amino acids and sulphur peptides represents a source of aromatic precursors that the yeast can use for the production of tropical aromas. Bouquet B49 helps to produce wines with stable, fresh and intense aromas.

Composition: blend of gallic tannins and inactive yeast rich in peptides.

Main application: production of white and rosé wines rich in tropical notes.

Dosage:

• rosé must: 5-20 g/hL.

white must: 2-20 g/hL.

We recommend combining each product in the range with its perfect companion yeast





SPECIAL TANNIN-BASED PRODUCTS FOR VINIFICATION

ESSENTIAL RANGE



The ESSENTIAL range of tannins is the result of a broad programme involving the selection of the best woods and essences and the study of the effects of the different extraction techniques under different working conditions. ESSENTIAL tannins also undergo stringent tests in order to ensure constant quality in all the batches produced. Within the ESSENTIAL range you can find the right tannin for every winemaking need and for every technical requirement. Furthermore, they come with advice on the right dosage for each different style of wine and for the various markets around the world. We always recommend performing a taste test before treatment.


0,5 kg


0,5 kg



ESSENTIAL PEP

Tannin extracted from grape pips





Composition: condensed tannins derived from grape pips.

Main applications: during the fermentation and racking of red and rosé must, for colour stabilization. During the aging and pre-bottling stages, to improve the body and the aromatic complexity of the wine.

Sensory characteristics: complexity on the nose, enhancement of varietal notes.

• red must and wine: 5-20 g/hL.

• white and rosé must and wine: 1-10 g/hL.

(see table at the end of this section)



ESSENTIAL PEL

Tannin extracted from grape skins

When used during the alcoholic fermentation, this tannin enhances the antioxidant power of the must and preserves its feeling of freshness. When used during the aging and the pre-bottling stages, it refreshes aro-

mas by enhancing their fruitiness, while increasing the structure and rebalancing the mouthfeel of the wine. Essential PEL is perfect for stepping up the quality of wines at the same time as increasing their aromatic intensity and softness.

Composition: tannin extracted from the skins of white-wine grapes.

Main applications: during the fermentation and racking of white, rosé and red must to increase antioxidant protection and create long-lasting freshness.

Sensory characteristics: freshness on the nose, rebalance of the mouthfeel.


• red must and wine: 5-20 g/hL. • white and rosé must: 1-10 g/hL. (see table at the end of this section)

TANNINS SPECIAL TANNIN-BASED PRODUCTS FOR VINIFICAT



ESSENTIAL OAK BARREL

Oak tannin



notes. Essential Oak Barrel can be used to supplement the loss of tannins from pre-used barrels, meaning that it can be used for longer without losing its ability to induce the development of the wine. Essential Oak Barrel combats reduction in wines aged in steel tanks or used casks.

Composition: ellagic tannin derived from oak wood.

Main applications: used during the pre-bottling stage for red wines, helps boost the structure and intensity of aromas as well as providing an antioxidant effect.

Sensory characteristics: vanilla and coconut.

• pre-bottling of white, rosé and red wines: 1-20 g/hL. (see table at the end of this section)



ESSENTIAL OAK SWEET

Blend of hydrolyzable tannins

Using Essential Oak Sweet enhances the sense of sweetness and intensity of aromas and brings wines back into balance when they've been aged in oak barrels.

Composition: Oak tannin.

Main applications: a useful tool for improving white, rosé and red wines during the pre-bottling stage, with an excellent antioxidant effect.

Sensory characteristics: coconut, sweetness.

Dosage:

• pre-bottling of white and rosé wines: 1-5 g/hL.

pre-bottling of white and rosé wines: 3-20 g/hL.

(see table at the end of this section)



















TANNINS SPECIAL TANNIN-BASED PRODUCTS FOR VINIFICATION

0,5 kg



ESSENTIAL FREE VEG

Blend of tannins extracted from the wood of exotic species



Using Essential Free Veg during the aging stage helps cover up vegetal and astringent notes, and allows the varietal properties which tend to hide behind 'green' notes to come to the fore.



0,5 kg

Composition: condensed tannins extracted from the wood of exotic species.

Main application: covers up vegetal notes when aging white, rosé and red wines, with an excellent antioxidant effect.

Sensory characteristics: helps cover up vegetal and astringent notes.


Dosage:

• for aging rosé and red wines: 5-20 g/hL. (see table at the end of this section)



ESSENTIAL FREE OFF

Oak tannin



responsible for off-odours in wines. Essential Free Off contains a high level of ellagitannins and simple phenols, which are very active as oxygen receptors. This makes it a valuable weapon in the fight against oxidation.

Composition: hydrolyzable tannin.

Main applications: used when aging white, red and rosé wines, it helps cover any notes of reduction and increases the wine's structure.

Sensory characteristics: solve the problems linked to the perception of reduction and draw out varietal notes.


Dosage:

• for aging white, rosé and red wines: 0,5-20 g/hL. (see table at the end of this section)



ESSENTIAL OAK STRONG

Blend of condensed and hydrolyzable tannins



been aged in oak barrels. Also enhances aging notes and tertiary aromas.

SPECIAL TANNIN-BASED PRODUCTS FOR VINIFICAT



TANNINS

0,25 kg

Composition: condensed and oak tannins.

Main applications: used during the pre-bottling stage for red wines, helps boost the structure and intensity of aromas as well as providing an antioxidant effect.

Sensory characteristics: liquorice, tobacco.

Dosage:

• pre-bottling of red wines: 3-20 g/hL. (see table at the end of this section)





ESSENTIAL PASSION

Blend of condensed tannins

Using Essential Passion when aging wines enhances the fruity notes in the wine, as well as the perception of sweetness and the intensity of varietal aromas.



Composition: condensed tannins extracted from the wood of red-berry trees.

Main application: when aging red and rosé wines, this product is helpful to boost the sensory profile as well as providing an antioxidant effect.

Sensory characteristics: sweetness, fruity notes, intensity of varietal aromas.

Dosage:

• for aging red wines: 10-20 g/hL. (see table at the end of this section)





TANNINS SPECIAL TANNIN-BASED PRODUCTS FOR VINIFICATION



ESSENTIAL OAK PROGRESS

Blend of condensed and hydrolyzable tannins





Composition: condensed and oak tannins.

Main applications: when aging white, rosé and red wines, this product is helpful to boost the complexity, structure and antioxidant effect.

Sensory characteristics: fruity notes, sweetness, persistence.

• aging of white and rosé wines: 1-5 g/hL.

• aging of white and rosé wines: 10-20 g/hL.

(see table at the end of this section)



ESSENTIAL FRESH

Blend of condensed and gallic tannins

Used at the aging and pre-bottling stages, Essential Fresh helps refresh tired, oxidized wines and open up their varietal notes. Using Essential Fresh balances the structure of wines by heightening the sensations of

structure that contribute significantly to the improvement of taste and retronasal perception. Wines treated with Essential Fresh display more complexity and persistence of aromas and flavours. It boosts the effects of SO₂ thanks to the blend of different types of tannins. Great at preserving the intensity of aromas and flavours in white wines.

Composition: blend of condensed and gallic tannins.

Main applications: when used during the aging and pre-bottling of white, rosé and red wines, it is useful to cover up notes of oxidation at the same time as increasing freshness and opening up aromas and flavours.

Sensory characteristics: freshness, complexity, persistence.

Dosage:

• at the aging and pre-bottling stages for white, rosé and red wines: 0.5–10 g/hL. (see table at the end of this section)

0,5 kg






0,5 kg



PRIVILÈGE BLEU

Blend of oak tannins



out its varietal notes and increasing its complexity. Privilège Bleu can also be used when aging wines to help prolong the lifespan of the wooden vessels.

Composition: blend of oak-wood tannins (from different species).

Main application: enhance the structure, fullness and intensity of aromas during the pre-bottling of red, white and rosé wines.

Sensory characteristics: coconut, chocolate, coffee.

Dosage:

• pre-bottling of white, red and rosé wines: 1-20 g/hL. (see table at the end of this section)



PRIVILÈGE NOIR

Blend of oak tannins

When used during the pre-bottling stage, Privilège Noir enhances the fruity notes arising from varietal characteristics, sensations of sweetness and fullness. It also helps balance wines when they've been aged in

oak barrels. For wines with an elegant, refined and austere sensory profile, using Privilège Noir provides the missing elements to achieve real quality.

Composition: blend of oak-wood tannins (from different species).

Main applications: enhance the fullness and intensity of aromas during the pre-bottling of red and rosé wines, as well as providing an antioxidant effect.

Sensory characteristics: red berry fruit, jam.

Dosage:

• pre-bottling of red and rosé wines: 1-20 g/hL. (see table at the end of this section)



SPECIAL TANNIN-BASED PRODUCTS FOR VINIFICAT

PRIVILEGE RANGE



Every tannin is unique in terms of both its chemical-physical properties and its interaction with the wine. The PRIVILÈGE range of tannins produced by the IOC group is the result of selection of the finest base materials, careful combination of these materials and an exclusive extraction process distinguished by low-pressure and low-temperature working conditions. These conditions make it possible to extract solely and exclusively the tannins that will give the wine added structure and softness, as well as enhancing the overall complexity of the wine by drawing out the unique characteristics of each variety of grape.

0,25 kg















SPECIAL TANNIN-BASED PRODUCTS FOR VINIFICATION

0,25 kg



PRIVILÈGE DUO

Blend of oak-wood tannins to use in malolactic fermentation



Privilège Duo is an oak-tannin-based preparation designed specifically to be used in synergy with the Maxiflore Satine bacterium 48hrs before inoculating, in order to enhance the aroma in red wines.




Composition: blend of oak-wood tannins (from different species).

Main application: enhance the aroma in red wines.

Dosage:

• 10 g/hL.



SPECIAL TANNIN-BASED PRODUCTS FOR VINIFICATION

Average doses in g/hL

Perform laboratory tests to establish the technical effects the tannin has on the sensory profile and determine the ideal dosage.

	T N	AUST	AG	ING	PRE-E	BOTTLING	CHARACTERISTICS
Essential PEP	1 - 10	5 - 20	1 - 10	5 - 20	1 - 10	5 - 20	Colour, aromatic complexity, structure, enhancement of varietal notes.
Essential PEL	1 - 10	5 - 20	1 - 10	5 - 20	1 - 10	5 - 20	Freshness on the nose, rebalance of the mouthfeel.
Essential OAK Barrel					1 - 5	5 - 20	Coconut, vanilla, structure, intensity of aromas.
Essential OAK Sweet			1 - 5	10 - 20	1 - 5	3 - 20	Coconut, sweetness, intensity of aromas.
Essential Free Veg			5 - 10	10 - 20			Helps cover up vegetal and astringent notes.
Essential Free Off		5 - 10	0.5 - 2	5 - 20			Solves the problems linked to the perception of reduction.
Essential OAK Strong			1 - 5	5 - 20	1 - 5	3 - 20	Liquorice, tobacco, structure, intensity of aromas and length.
Essential Passion			1 - 5	10 - 20			Sweetness, fruity notes, intensity of varietal aromas.
Essential OAK Progress			1 - 5	10 - 20			Fruity notes, sweetness and persistence.
Essential Fresh					0.5 - 10	1 - 10	Freshness, complexity, persistence.
Privilège Bleu					1 - 2	1 - 20	Coconut, chocolate, coffee, structure, fullness, intensity of aromas.
Privilège Noir					1 - 2	1 - 20	Red fruit, jam, fullness, intensity of aromas.

^{*} if made from Botrytis-infected grapes: 5–10 g/hL, on finished wine: 1–4 /hL





In compliance with OIV resolution OENO 430/2010, our woods do not undergo any form of combustion – not even on the surface – and they aren't charred or crumbly to the touch. Moreover, they have not undergone any form of chemical, enzymatic or physical treatment except heating through convection.

	Sensory impact	(eg)wood!	্রি Dosage g/hL	Different types of chips	Average time of contact
	first-use barrel effect	FEEL WOOD PREMIUM BARREL R	300-600	2	25-60 days
	used barrel effect	FEEL WOOD SHADE BARREL R	300-500	2	25-60 days
	enhancement of red-berry notes	FEEL WOOD RED FRUITS R	300-500	2	25-60 days
*	sensory complexity and spicy notes	FEEL WOOD SPICY FLAVOR R	150-300	3	25-60 days
D	sensations of chocolate and cappuccino	FEEL WOOD DARK CHOC R	150-300	1	25-60 days
	used barrel effect	FEEL WOOD SHADE BARREL W	100-200	2	25-60 days
Ô	sweet and fruity notes	FEEL WOOD WHITE FRUITS W	50-150	1	25-60 days
	floral aromatic notes	FEEL WOOD SENSE FLORAL W	100-200	2	25-60 days

WOODEN BARREL SUBSTITUTES

FEELWOOD! RANGE

As the conclusion to a research programme carried out over several years, the IOC group has selected a special range of high-quality wood derivatives from a variety of different sources and with different levels of toasting, the aim being to give each product in this family a specific, unique sensory objective.

The study was conducted by testing different combinations of these base ingredients on many different varieties of wine until a complete range for both white and red wines was created. FeelWood gets its name because the "feel" produced by each product in the range is the winemaker's main aim in using wood derivatives to craft their wine.

The wood chips are packaged in specially designed infusion bags. The bags have a divider keeping the various types of wood separate to ensure the homogeneity of the formulations. This homogeneity wouldn't be guaranteed by mixing up different sizes of chips. Chip sizes: 0.5-2cm.



10 kg



PREMIUM BARREL R

Chips in Infusion Bag

Composition: blend of selected French and American species of oak (*Quercus sp.*).

Main applications: closely imitate the sensory effects obtained by using a new oak barrel. To be used on red wines.

Dosage:

• 300-600 g/hL for 25-60 days.





SHADE BARREL R

Chips in Infusion Bag

Composition: blend of selected American species of oak (Quercus sp.).

Main application: replicate the sensory effect of used oak barrels for aging

red wines.

Dosage:

• 300-500 g/hL for 25-60 days.



Chips in Infusion Bag



10 kg

10 kg

10 kg

10 kg

Composition: blend of selected French species of oak (Quercus sp.).

Main applications: enhance and amplify notes of red berries in the bouquet

of red wines.

Dosage:

• 300-500 g/hL for 25-60 days.



SPICY FLAVOR R

Chips in Infusion Bag

Composition: blend of selected French species of oak (Quercus sp.).

Main applications: enhance and amplify aromatic complexity in general, as well as spicy notes. To be used on red wines.

Dosage:

• 150-300 g/hL for 25-60 days.



DARK CHOC R

Chips in Infusion Bag

Composition: blend of selected French species of oak (Quercus sp.).

Main applications: provide an intense chocolately feel, with added notes of coffee, hints of spice and a touch of coconut. To be used on red wines.

Dosage:

• 150-300 g/hL for 25-60 days.



SHADE BARREL W

Chips in Infusion Bag

Composition: blend of selected American species of oak (Quercus sp.).

Main applications: enhance and amplify the best aromatic notes connected with aging white wines in used barrels.

Dosage:

• 100-200 g/hL for 25-60 days.

WOODEN BARREL SUBSTITUTES FEELWOOD! RANGE



WHITE FRUITS W

Chips in Infusion Bag

Composition: blend of selected American species of oak (Quercus sp.).

Main applications: enhance and amplify the fruity notes in white wines.

Dosage:

• 50-150 g/hL for 25-60 days.



SENSE FLORAL W

Chips in Infusion Bag

Composition: blend of selected American species of oak (Quercus sp.).

Main applications: enhance and amplify the floral notes in white wines.

Dosage:

• 100-200 g/hL for 25-60 days.





10 kg





Filter surfaces aren't always able to capture all the suspended particles in a wine, especially if the liquid being filtered is rather diluted. Filter aids are very useful, as they increase the level of filtration and reduce clogging.

The Perdomini-IOC range comprises three types of filter aid:

- Perlites
- Fossil shell flours
- Precoat filtration

PERLITES

25 kg - 22 kg 20 kg



Filter aid

DIAPERL

The perlite products in the Diaperl range display three different levels of permeability, as can be seen from the table below. The main features of these perlites are that they are ultra-pure and free from organic residue – this means they create no unwanted odours in the filtered product.



Composition: perlite.

Main application: form filtration cakes in DE and vacuum filtration.

Dosage: pour Diaperl into the product being filtered at a dosage ranging from 50-120g/hl, depending on specific needs. Excellent results have been attained by using Diaperl in tandem with Drenopor and Drenopor CF precoats.

ANALYTICAL DATA	DIAPERL M	DIAPERL V	DIAPERL VV
Permeability in Darcy	0,9	1,5	2,03
Wet specific weight (g/l)	180	150	150
Colour	white	white	white
Moisture	1%	1%	1%
Silica (SiO ₂)	72,0%	72,0%	72,0%
Alumina (Al ₂ O ₃)	14,0%	14,0%	14,0%
Iron oxide (Fe ₂ 0 ₃)	0,7%	0,7%	0,7%
Na ₂ O	4,0%	4,0%	4,0%
K ₂ 0	8,8%	8,8%	8,8%
Calcium oxide (CaO)	0,3%	0,3%	0,3%
Magnesium oxide (MgO)	0,1%	0,1%	0,1%

FILTRATION ADDITIVES

FOSSIL SHELL FLOURS

20 kg

DIACEL



Inert, insoluble and presenting a volume of distribution only 15% of which is constituted by solids, these filter aids are perfect for forming filter cakes for use in DE filtration.



Composition: top-quality American diatoms.

Main applications: primary filtration and polishing.

Dosage: to form the precoat, pour Diacel into water or another clear liquid at a dosage of 500-1000g/m² of filter surface. Excellent results can be attained by combining Diacel products with Drenopor and Drenopor CF precoats. When ready for filtration, pour 50-120g/hl (depending on specific needs) of Diacel into the product being filtered.

PRECOAT FILTRATION

10 kg

DIAPOL SUPER MV

Stabilizing filter aid



The Diapol Super MV formulation optimizes the respective proportions of PVPP, silica gel and inert substances. This product assists the filtration process and at the same time provides a powerful stabilizing effect on any unstable polyphenols and proteins in the wine. Removing proteins with silica gel leads to wines with a low filterability index, therefore with less risk of clogging up the microfiltration membrane.

Composition: PVPP, silica gel, fossil shell flours.

Main applications: lower the catechin and leucoanthocyanin content, stabilize wines affected by oxidation or protein clouding, stabilize colouring substances.

Dosage:

• 20–50 g/hL depending on the type of wine and the operating conditions.



FILTRATION ADDITIVES

DIAPOL VB

Stabilizing filter aid

Diapol VB is intended for exclusive use with white must and wine, to remove excess colour and unstable polyphenolic fractions, which are responsible for browning.

Diapol VB can be used directly in DE filtration, but acts more forcibly when added to the bulk for tank filtration. The removal of protein fractions by silica gel makes it possible to obtain wines with a low filterability index and therefore with less danger of clogging the microfiltration membrane.

Composition: PVPP, activated carbon, silica gel, fossil shell flours.

Main applications: helps carry out both the DE filtration process and an effective stabilizing action on the product.

Dosage:

• 60-120 g/hL depending on the type of wine and the operating conditions.

10 kg

10 kg

DRENOPOR CF 300

Precoats with strong adsorption capacity

Top-quality perlites combined with α -cellulose fibres for highly effective precoats. Cellulose performs a mechanical action forming a compact, easily removed precoat, while perlites form a complex network of tiny ducts that guarantee retention of the solid particles and other impurities. The precoat is also distinguished by its exceptional strength, which makes it resistant to breaking and cracking.



Composition: α-cellulose fibres, fossil shell flours.

Main application: Drenopor CF 300 is particularly recommended for high-yield filtration.

Dosage:

- 500-800 g/m² of filter surface.
- 20-50 g/hL for DE filtration.

0 kg

DRENOPOR FILTER "M" E "S"

Precoats with strong adsorption capacity

The synergy between the Drenopor Filter components leads to the formation of a precoat whose surface is perfectly homogeneous and free from preferential flow paths that could compromise the outcome of the filtration process. The precoat is also distinguished by its exceptional strength, which protects it from breaking and cracking.



Composition: α-cellulose fibres, fossil shell flours, perlites.

Main application: form precoats for DE filtration.

- DRENOPOR M is particularly recommended for high-yield filtration.
- DRENOPOR S can be used for "polishing" filtration.

Dosage:

- 400-700 g/m² of filter surface.
- 20-50 g/hL for DE filtration.

HYDROPOR

Prehydrated filter aid

The raw material used is subjected to a special mechanical process in order to obtain fibres that are more open, with a consequent increase in the relevant surface area, improved grip and the opportunity to make prepanels with uniformity of distribution (isotropic structure). Depending on the dosage and the permeability of the adjuvant used, Hydropor can be used both for coarse and polishing filtration. However, for particularly wide filtration, the use of Drenopor Filter M is more suitable in order to prevent clogging phenomena.

Composizione: α -cellulose fibres.

Main application: ideal for deep and DE filtration.

Dosage:

• 500-1000 g/m² of surface area.

HYDROPOR BLEND

Prehydrated filter aid

The α -cellulose fibres have been processed using a mechanical procedure that opens them up to the right point; this way, they ensure that the precoats adhere perfectly to the surfaces of the filter sheets without creating preferential flow paths. The diatoms used together with these fibres help disperse the fibres quickly and create the right degree of drainage to avoid sudden clogging problems.

Composition: α-cellulose fibres, diatoms.

Main applications: DE filtration of must and wine. Suitable principally for polishing processes.

Dosage:

• 500-1000 g/m² of surface area.

PRECOAT

Prehydrated filtration adjuvant

The products in the Precoat range contain alpha-cellulose fibres processed using a special procedure that guarantees both optimal filtration and ease of use.

- Precoat SV Wet density: 0,268 g/mL Permeability: $300-500 \, hL/h/m^2$
- Precoat M Wet density: 0,266 g/mL Permeability: 250-500 hL/h/ m²
- Precoat S Wet density: 0,298 g/mL Permeability: 200-350 hL/h/ m²

Composition: precoat with medium α -cellulose-fibre content and silicates.

Main application: DE filtration, from primary filtration to polishing.

- Precoat SV: for high-yield DE filtration and rotary vacuum filters. Used to filter must, juice and liquids with dregs.
- Precoat M: particularly recommended for high-yield DE filtration;
- Precoat S: recommended for polishing filtration.

Dosage:

- Precoat SV: 10% of the weight of filtration flour to be used.
- Precoat M and S: 400–700 g/m² of filter surface. 20–100 g/hL for DE filtration.





















Our Perdomini-IOC range of cleaning products is backed up by a full assistance and advice service which not only identifies the most suitable products for each individual need, but also provides assistance with using the products, training for the staff who will use them and drawing up hygiene protocols in compliance with the most recent legislation (HACCP, safety and environment).

To draw up a suitable, detailed operations protocol, we advise you to contact the Perdomimi-IOC technical assistance service.



ACIDIC DETERGENTS

10 kg - 25 kg 250 kg

DETAR 10

Surfactant-free acidic descaling detergent

Use at % concentration: 1-3

Characteristics and main applications: solubilisation and removal of precipitates - whether inorganic and calcareous or mixed (organic and inorganic) - from the surfaces of machines, plants and separation systems.

Detar 10 is the result of development aimed at managing membrane separation systems, completing the primary cleaning process for organic material. Above all in cross-flow systems, which are required to operate in wildly differing conditions and be regenerated with water of varying hardness and composition, Detar 10 has demonstrated that it contributes significantly to the maintenance of the overall performance of the system over time.

DETAR ACID

Acid detergent and descaling agent with surfactant

Use at % concentration: 2-5

Characteristics and main applications: detergent designed for removing inorganic deposits from the surfaces of machines and equipment. Detar Acid has been specially formulated to dissolve and remove inorganic residues, whether caused by processing operations or the utility water supply (limescale). Detar Acid's formula has been optimized through the use of surfactants and corrosion-inhibiting substances. Detar Acid produces only a limited quantity of foam and can therefore be used for circulation cleaning.

> 25 kg 1100 kg

DETAR ACID ASF

Acid detergent and descaling agent

Use at % concentration: 0.5-5

Characteristics and main applications: break up and remove limescale deposits and eliminate mixed residues (organic and inorganic) from tanks, pipes, pasteurizers and heat exchangers. Detar Acid ASF is particularly effective as a follow-up to alkaline cleaning operations to provide complete neutralization of surfaces and equipment.







Acid detergent and lime scale remover with a sanitizing action

Use at % concentration: 3-5

Characteristics and main applications: Detar Acid San is a more than valid alternative to traditional cleaning processes that involve initial cleaning with an alkaline product, followed by an acid and finally a sanitizing treatment, of the inside of vats, filling machines, piping or storage tanks and pasteurizers. The Detar Acid San formula has been expressly designed to avoid foam production, making this product particularly suitable for CIP and water recirculation systems.

> 25 kg 1100 kg

DETAR SAN SCHIUMA

Acidic foaming detergent and lime scale remover with added sanitizing power

Use at % concentration: 4-8

Characteristics and main applications: with the use of special equipment, the unique formulation of Detar San Schiuma leads to the production of a dense, clingy foam that makes it easy to clean vertical surfaces and walls. Also recommended to clean walls, floors and the outside of storage tanks and vats.

ALKALINE DETERGENTS

1300 kg

DETAR FLOW

Highly alkaline detergent

Use at % concentration: 1-3

Characteristics and main applications: Thanks to its power to sequester and suspend solid particles, Detar Flow is particularly effective at removing large quantities of organic contaminants from both winemaking equipment and filter membranes. Suitable for removing the high concentration of organic substances left behind in cross-flow filtration systems, especially at the final stage of the processes when the organic elements have been separated and concentrated, and form layers which stubbornly resist removal. Detar Flow has been shown to make a measurable contribution to stabilization and to the long-term continuation of filtration-system performance.

25 kg 1000 kg



DETERGENTS ALKALINE DETERGENTS

1300 kg

DETAR MEM

EDTA-free highly alkaline detergent

Use at % concentration: 1-3

Characteristics and main applications: thanks to its sequestering action, Detar MEM is particularly effective at removing large quantities of organic fouling. As it does not contain EDTA, this product is highly suitable for use in waste water treatment and purification plants. Detar MeM is recommended for cleaning containers, tanks, pipes and pasteurizing lines using either automatic or CIP cleaning systems.

> 10 kg 25 kg

DETAR PS

Alkaline detergent with surfactant

Use at % concentration: 2-4

Characteristics and main applications: Detar PS is a product formulated specifically for the purpose of optimizing operations to regenerate cartridges and filter membranes when simple physical regeneration with water and/or steam has failed to achieve the desired results.

> 25 kg 1100 kg

DETAR SCHIUMA

Foaming alkaline detergent

Use at % concentration: 4-6

Characteristics and main applications: Medium-alkalinity foaming detergent suitable for removing stubborn organic residues (both animal and non-animal fats) from surfaces. The use of Detar Schiuma, a foaming detergent, makes it possible to carry out cleaning operations on machinery, equipment and walls where there are vertical and diagonal surfaces or any other surfaces hard to clean using traditional methods.

> 1 kg - 25 kg 1400 kg

DETAR TRE LIQUID

Caustic descaling detergent

Use at % concentration: 0.5-2

Characteristics and main applications: Detar Tre Liquid posis a highly caustic detergent, which means it has powerful degreasing and tartrate-removal effects. The inclusion of various kinds of organic sequestering agents in the formula gives this product a strong complexing action. This feature means that Detar Tre Liquid can also be used where the water is particularly hard.









Caustic detartrating detergent in powder form

Use at % concentration: 5-10

Characteristics and main applications: complete detergent in powder form, highly alkaline and phosphate free. Detar Tre Special is perfect for cleaning and removing tartrates from stainless steel equipment such as vats and conveyors, as well as pasteurizers, concentrators and centrifuges. It's also suitable for use on presses and filters. This product is compatible with light allovs, zinc and aluminium.

PERLAC CF/O

Powder detergent for bottle washers

Use at % concentration: 5-10

Characteristics and main applications: Perlac CF/O has been specially designed for use in bottle washers. This all-in-one formula fully satisfies all the different demands of bottle cleaning and preparation for reuse thanks to the special surfactants included in its formula. While performing its cleaning and unsticking actions, Perlac CF/O does not foam, thanks to the careful choice of non-foaming surfactants.

PERLAC CF/3

Powder detergent for bottle washers

Use at % concentration: 1-3

Characteristics and main applications: Perlac CF/3 is a complete product, rich in sequestering agents and with special highly biodegradable surfactants working in synergy with the alkaline salts to guarantee great performance every time. Perlac CF/3 has been specially designed for use in bottle washers and is particularly recommended when the water supply is especially hard. The synergic action of the alkaline environment carefully balanced with dispersing agents and surfactants allows labels to be peeled off whole, thus avoiding the problem of pumps and filters becoming clogged up. While performing its cleaning and unsticking actions, Perlac CF/3 does not foam, thanks to the careful choice of non-foaming surfactants, which best display this characteristic at temperatures of around 50-55°C.

DETERGENTS ALKALINE DETERGENTS

1400 kg

PERLAC CF/LIQUID

25 kg

25 kg

Caustic detergent, phosphorus-free

Use at % concentration: 1-3

Characteristics and main applications: Perlac CF/Liquid guarantees complete removal of processing residues from conveyor belts, bottling plants, pasteurizers and tanks. Its composition has been carefully studied in order to create a product that would simultaneously meet the cleaning needs of the food industry and fulfill the ever stricter requirements of the environment sector. It is a caustic product with added surfactants and sequestering agents with performance levels that allow it to be used effectively even in limit conditions (e.g. particularly hard water, even above 25°F; temperature higher than 70°C). It is also suitable for removing tartrate deposits from containers and concentrators and for cleaning floors and walls that are particularly dirty with organic matter.

PERLAC FILL

Highly caustic detergent

Use at % concentration: 1.5-5

Characteristics and main applications: Perlac Fill is a product which stands out not only for its highly caustic effect, but also for its surfactants and sequestering agents. The surfactants in its formula stop it from foaming at a wide range of temperatures and independently of the type of dirt being treated. The detergent effect of the alkaline salts is also boosted by a powerful emulsifying and suspending action. Whether in CIP or circulation cleaning, Perlac Fill can be used to clean storage tanks, pasteurizers, fillers and piping. Excellent results can be obtained when cleaning floors and walls by hand, especially anywhere there's a lot of organic dirt.

PERLAC TENDER

Alkaline detergent

Use at % concentration: 3-5

Characteristics and main applications: Perlac Tender is recommended for cleaning containers, tanks, piping and pasteurization lines, whether CIP or automatic cleaning systems are used. It's particularly effective when combined with Detar 8. It's perfect for cleaning containers, piping, tanks, bottles, pasteurizers, fillers, crate washers, etc.













CHLORINATED ALKALINE DETERGENTS

10 kg

PERLAC CLORAT PRESSE

Chlorinated alkaline sanitizer

Use at % concentration: 0.5-2

Characteristics and main applications: Perlac Clorat presse is a sanitizing detergent containing surfactants, with high sequestering power. Phosphate free. As it frees chlorine, with Perlac Clorat Presse you can both clean and sanitize in a single operation. It is particularly recommended for washing, sanitizing and removing tartrate residues from presses with stainless steel cages. It can also be used to good effect to treat conveyor belts, vats and storage tanks. When used with a high-pressure hose, it foams slightly.

25 kg

PERLAC DEGREASER

Concentrated alkaline detergent

Use at % concentration: 0.5-2

Characteristics and main applications: Perlac Degreaser is a liquid alkaline detergent rich in active substances that is suitable for the removal of particularly heavy dirt from floors and walls. The simultaneous presence of alkaline substances, chelating agents and a pool of special surfactants ensures an incisive effect on the most varied types of dirt (animal and vegetable fats, oils, dust). The chelating substances contained in Perlac Degreaser result in the use of water with high hardness as well, for the dilution of the product.

> 25 kg 1200 kg

PERLAC SAN

Chlorinated alkaline sanitizer

Use at % concentration: 0.5-1

Characteristics and main applications: phosphate-free, chlorinated alkaline sanitizer. Its high chlorine content means that excellent cleaning and sanitizing results can be attained from a single application. Perlac San can be used to clean, sanitize and remove staining from wooden barrels and from storage and fermentation tanks made of both cement and stainless steel. Perlac San is also effective when used to sanitize floors and walls. It gets rid of unpleasant odours emanating from drainage channels and gutters. It can be used as a sanitization additive in the penultimate rinse phase in bottle-washing machines.

DETERGENTS CHLORINATED ALKALINE DETERGENTS

PERLAC SAN SCHIUMA NEW

Foaming chlorinated alkaline detergent

Use at % concentration: 3-5

Characteristics and main applications: Perlac San Schiuma New is particularly effective at removing organic residues and mould. Perlac San Schiuma New is recommended in any situation where there is a need for cleaning and contemporaneous sanitization in environments and on apparatus fouled with proteins and grease. When used with a special jet pipe, the foaming agents contained in this formulation generate an exceptionally thick, sticky foam that, when sprayed on vertical walls, leads to an increased contact time with the dirt. The result is effective cleaning and sanitization in one simple operation.

> 10 kg - 25 kg 200 kg - 1100 kg

PERLAC SAN SPECIAL

Chlorinated alkaline sanitizer

Use at % concentration: 0.5-1

Characteristics and main applications: chlorinated alkaline sanitizer with excellent cleaning properties. Phosphate free. Its special sequestering agents mean that excellent cleaning and sanitizing results can be attained, even when the water supply is especially hard. Perlac San Special is ideal for cleaning conveyor belts, vats, storage and fermentation tanks, and for eliminating the unpleasant odours that can form in drainage channels and gutters. Perlac San Special is non-foaming, so it can be easily used in CIP systems.









10 kg 25 kg

DETAROX

Sanitizer containing peracetic acid

Use at % concentration: 0.02-0.5

Characteristics and main applications: balanced blend with an oxidating effect, containing stabilized peracetic acid and hydrogen peroxide. Its ease of rinsing and super-low foam production make this product perfect for use in filtration systems. Detarox is recommended for cleaning and removing colour staining from fermentation tanks and bottling lines. Its superlow foam production makes Detarox perfect for use in all CIP systems. It doesn't present any risk of corrosion on stainless steel, aluminium, tin-plated iron or plastics (polyethylene, for example). Lastly, the unique composition of Detarox allows it to be used at low temperatures with guaranteed great performances.

> 10 kg 20 kg

DETAROX AP

Highly effective broad-spectrum bactericidal disinfectant

Use at % concentration: 0.2-0.5

Characteristics and main applications: highly effective broad-spectrum bactericidal disinfectant containing stabilized peracetic acid, formerly classified as a medico-surgical material, now Biocide. Detarox AP is non-corrosive on a wide range of materials including stainless steel, aluminium and many kinds of plastic. It is particularly recommended for systems, surfaces or packing whose level of hygiene can influence the finished product either directly or indirectly. Its wide-ranging action spectrum, easy-rinse formula and low foaminess make it perfect for the sanitization of filtration systems.

750 ml

DETAR SPRAY S

Alcohol spray sanitizer

Use at % concentration: ready for use.

Characteristics and main applications: Detar Spray S is a ready-to-use, alcohol-based cleaner and sanitizer, suitable for cleaning surfaces and equipment used for food processing, such as conveyor belts, cappers, taps, etc., in any sector of the food industry where the proliferation of microorganisms can constitute a serious problem. This product is perfume-free and colourant-free and is compatible with all surfaces in steel, plastic and aluminium. It is a dual-action degreaser and sanitizer.



DETERGENTS FCO-FRIENDLY SANITIZING DETERGENTS



ECO-FRIENDLY SANITIZING DETERGENTS

25 kg 1400 kg



DECO FILL

Caustic detergent free from E.D.T.A. and phosphorus

Use at % concentration: 1.5-5

Characteristics and main applications: Deco Fill does not contain phosphates or phosphonates and is completely free from phosphorus. No EDTA/NTA. Highly caustic detergent designed for use in the food and beverage industry. Deco Fill is a product which stands out not only for its highly caustic effect, but also for its surfactants and sequestering agents. Deco Fill can be used for CIP operations on processing and filling circuits.

> 25 kg 1300 kg



DECO FLOW

Caustic detergent free from E.D.T.A. and phosphorus

Use at % concentration: 1-3

Characteristics and main applications: DecoFlow does not contain phosphates or phosphonates and is completely free from phosphorus. No EDTA/NTA. DecoFlow has been developed specifically for particularly tricky operations to regenerate cross-flow filtration systems and microfiltration systems. Under normal conditions, displays exceptional compatibility with metals - steel above all - and with a very broad range of polymer and non-polymer filter membranes.









DECO PS

Caustic detergent free from E.D.T.A. and phosphorus

Use at % concentration: 2-4



25 kg 1100 kg

Characteristics and main applications: Deco PS is a product formulated specifically to optimize filter cartridge and membrane regeneration operations whenever simple physical regeneration with water and/or steam is insufficient. Deco PS removes the organic residue clogging the filters which has accumulated during the filtering process. In particularly difficult cases to resolve – for example when faced with stubborn, sticky matter – it may be helpful to use Deco PS and Detar 8 together. This way the alkaline and oxidizing action is backed up by powerful mechanical dirt removal.

CLEANING FILTRATION EQUIPMENT AND SYSTEMS

TITAN I

Additive for cleaning cross-flow filtration systems and apparatus

Characteristics and main applications: formulation containing dispersants and chelating agents suitable for the removal of mineral/inorganic residues

(manganese, iron, silica, aluminium, magnesium and calcium) primarily caused by the water supply, which deposits these residues inside piping and filtration apparatus, causing crossflow filtration systems to clog up. When used after treatment with Titan O (removal of organic materials), it helps maintain efficient performance in the cross-flow filtration systems used in wineries.

TITAN O

Enzyme-based product to enhance cleaning operations on cross-flow filtration systems and apparatus

Characteristics and main applications: Titan O is a powerful, non-foaming cocktail of enzymes that enhances the cleaning of organic residues (proteins, cellulose, starch, organic colloids from the wine, etc.) from the cross-flow filtration systems and apparatus used in wineries. Thanks to its mixture of enzymes, Titan O ensures deep but non-invasive cleaning of materials and is particularly effective at removing the red pigments and tannins found in red wines. Titan O has zero environmental impact (made of 100% biodegradable natural enzymes) and, when used routinely, it maintains optimal performance levels of filtration.




OTHER PRODUCTS

25 kg

ANTISCHIUMA

Silicon-based defoamer



10 kg - 25 kg 200 kg - 1100 kg

DETAR 8

Peroxide-based detergent

Use at % concentration: 1-2

Characteristics and main applications: specially designed for the removal of particularly difficult food residue. Thanks to the lack of foaming and the presence of sequestering agents, the best use for Detar 8 is in CIP systems, always in conjunction with an alkaline product such as Detar PS. The effectiveness of Detar 8 is perfectly exploited in the delicate operation of periodically regenerating filter cartridges.

PERLAC NASTRI

Lubricant for conveyor belts

Use at % concentration: 0.8-2

Characteristics and main applications: this lubricant for conveyor belts performs both a detergent and a sequestering action. It's a crucial aid to keeping bottling and packaging lines running smoothly. Perlac Nastri has been specially formulated in order to help conveyor belts run more smoothly. As well as improving productivity, Perlac Nastri also keeps the various components of conveyor belt systems in good working order for longer. Thanks to its special formula complete with surfactants, Perlac Nastri adds top cleaning power to its crucial role as a lubricant, thus ensuring the emulsification of any residues left behind by accidental bottle breakages.






DETERGENTS OTHER PRODUCTS

PERLAC NASTRI PLUS

Lubricant for conveyor belts

Use at % concentration: 0.2-0.4



25 kg - 200 kg 1000 kg

PERLAC NASTRI SPECIAL

Lubricant for conveyor belts

Use at % concentration: 0.2-0.4

Characteristics and main applications: pH-neutral belt lubricant with added cleaning and sanitizing power. Perlac Nastri Special contains special substances which work together to form a thin and effective lubricating layer between the conveyor belt and the bottles. This drastically reduces the amount of friction, consequently improving the efficiency of the conveyor, with fewer bottles tipping over or becoming stuck because of the belt moving jerkily. Moreover, thanks to this product the belt stays intact and in good working order for longer. Perlac Nastri Special performs a sanitizing action.

25 kg

PERLAC NEUTRO

Detergent with surfactants for general use

Characteristics and main applications: detergent based on efficacious vegetable-origin solubility enhancers acting in synergy with surfactants, emulsifiers and dispersing agents in a pH-neutral formula. With Perlac Neutro you can be sure your surfaces won't suffer from corrosion and your cleaning staff won't be exposed to hazardous chemicals. It's also an extremely versatile product, as it can be used for cleaning with both manually operated apparatus and automatic devices. Perlac Neutro can be used to clean the floors in warehouses and processing rooms – especially those where wheeled vehicles pass through – as well as offices and laboratories.

200 kg











completeran



Perdomini-IOC offers a complete range of products for the various stages of winemaking, including yeasts, fermentation starters, bioprotectors, stabilizers, fining agents, enzymes, tannins and alternatives to wooden barrels.



All our products have been formulated as the result of targeted experimentation programmes and each one arrives brings its own unique philosophy to your winery.



Each product that makes it through the selection process is further tested both in-house and at independent laboratories or testing agencies so that our customers can be sure what they're buying is reliable, compliant and safe for end consumers.



By blending careful selection of ingredients, preparation of the formulas following precise protocols, scrupulous hygiene practices and care for the environment, Perdomini-IOC is a guarantee of quality products in your winery.









Perdomini-IOC S.p.A.

Via Salvo D'Acquisto, 2 – 37036 San Martino B.A. (VR) Italy Tel. +39 045 8788611 r.a. info@perdomini-ioc.com www.perdomini-ioc.com

Inspired by nature.





